



## Comparative effectiveness of certain insecticidal compounds on the development and adult performance of red palm weevil, *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae)

Tanani MA

Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Madenit Nasr, Cairo, Egypt

### Abstract

The red palm weevil *Rhynchophorus ferrugineus* is a serious pest of date palm in different regions of Middle East. The present study was carried out to evaluate comparatively the effects of pyriproxyfen (insect growth regulator), neemazal (plant product) and spinetoram (bacterial product) on growth, development and adult performance of this insect pest. The early 5th instar larvae were treated with five sublethal concentrations of each compound. The most important results can be summarized as follows. Both larval duration and weight were considerably reduced by Spinetoram. Pyriproxyfen significantly reduced the larval duration only at 4000 ppm ( $7.2 \pm 2.97$ , compared to  $11.7 \pm 1.6$  days control insects) but Neemazal could not exhibit a remarkable effect. The strongest reducing effect on larval weight was recorded for Pyriproxyfen, since the growth inhibition% was determined as 49.10, 48.58 and 38.04, at 4000, 2000 and 1000 ppm, respectively. Pupation was inhibited by all tested insecticidal compounds and the pupal duration was significantly shortened, except at the lower three concentrations of Neemazal. Some deformations had been recorded only in adults, irrespective of the tested compound. The adult longevity was significantly shortened at the higher concentrations of each compound.

**Keywords:** adult, development, growth, longevity, morphogenesis, neemazal, mortality, pyriproxyfen, Spinetoram, pupa

### 1. Introduction

The red palm weevil (RPW) *Rhynchophorus ferrugineus* Olivier is a dangerous insect pest of date palm in Arabian Gulf region since the mid-1980s and Egypt since 1992 (Saleh, 1992; Cox, 1993) [1, 2]. In almost all Egyptian governorates, the larval and adult stages existed around the year seasons (Helal and El-Sebay, 1994; El-Sebay, 2004; Merghem, 2011; Merghem and Bibers, 2014) [3, 4, 5, 6]. Serious damage has been recorded for the larval stage of this weevil in some palm genera and, thus, control operations should be carried out (EPPO 2008) [7]. The chemical control of RPW with conventional insecticides usually exhibits a limited success in addition to the insecticide resistance of palms as well as their chemical residues accumulated in the nature as pollutants. Also, the extensive use of these insecticides caused various hazards to environment, mammals, beneficial animals and arthropods (Abo El-saad *et al.*, 2001; Ranasinghe *et al.*, 2003; Gloria *et al.*, 2008; Koul and Walia, 2009) [8, 9, 10, 11].

Therefore, different insecticide alternatives should be taken in consideration. Some of these alternatives are insect growth regulators (IGRs), like Pyriproxyfen, plant extracts, like neem formulations, and microbial products, like Spinetoram. Pyriproxyfen (IGR) exerts its toxic effect through the disturbance of the normal endocrine functions in insects (Smet *et al.*, 1990; Oberlander *et al.*, 1997) [12, 13]. Plant extracts, such as neem seed and leaf extracts, as well as, azadirachtin formulations, like neemazal, are effective agents as antifeedant, repellents, growth disruptors, moulting inhibitors and oogenesis suppressors (Rembold and Sieber, 1981; Garcia and Rembold, 1984; Dorn *et al.*, 1986; Richer *et al.*, 1997; Ghoneim *et al.*, 2000) [14, 15, 16, 17, 18]. In a previous study, pyriproxyfen, neemazal and spinetoram were recorded as active agents causing mortality in RPW (Hamadah and Tanani, 2014) [19]. Also, Hamadah and Tanani (2017) [20]

determined the deleterious effects of these compounds on haemogram of the same weevil. Spinetoram (microbial product) has been derived from soil dwelling bacteria *Saccharopolyspora spinosa* and exerts its toxic action by contact or ingestion (Sparks *et al.*, 1998) [21]. It targeting a nicotinic acetylcholine receptor as well as  $\gamma$ -aminobutyric acid (GABA) gated chloride channels causes insect paralysis (Watson 2001; Sparks, 2004; Sarfraz *et al.*, 2005) [22, 23, 24]. Objective of the current study was to evaluate the disruptive effects of certain insecticide alternatives (pyriproxyfen, neemazal and spinetoram) on the growth, development and adult performance of *Rh. ferrugineus*.

### Materials and Methods

#### 1. Experimental Insect

A culture of the red palm weevil *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) was initiated by a sample of field-collected pupae. The culture was established at Faculty of Science, Al-Azhar University, Cairo, Egypt, on sugarcane stem under constant laboratory conditions ( $29 \pm 2^\circ\text{C}$ , 60-70 % RH, the light intensity of about 30 foot-candles) (Rahalkar *et al.*, 1972; Ranavare *et al.*, 1975) [25, 26].

#### 2. Tested compounds and larval treatments

Spinetoram 5% (radiant), Pyriproxyfen 10% (admiral) and Neemazal (azadirachtin 20%) were kindly obtained from the Central Laboratory of Insecticides, Agricultural Research Centre, Doqqi, Giza, Egypt. Each compound was dissolved in distilled water for preparing five concentration levels: 40000, 20000, 10000, 5000 and 2500 ppm for Pyriproxyfen; 4000, 2000, 1000, 500 and 250 ppm for Neemazal and 200, 100, 50, 25 and 12.5 ppm for Spinetoram.

The newly moulted last instar larvae (5<sup>th</sup>) of *Rh. ferrugineus* were fed (for 24h) on fresh pieces of sugarcane stem after dipping for ten minutes in different concentrations of each

compound but the control congeners were dipped in water only for equal time period. All sugarcane pieces were allowed to dry before offering to the larvae. A day after treatment, all larvae (treated and control) were provided with untreated piece of sugarcane. Three replicates of 10 individual of larvae per each were used for each concentration. Each individual larva was isolated in a suitable glass vial. All vials tightly cover and located in a large cage supplied with a suitable electric bulb. At one day after feeding, the larvae were carefully weighted every day and also examined for mortality or other criteria.

### 3. Criteria of study

- **Body weight:** The fresh body weight was obtained every day using a digital balance.
- **Growth inhibition:** It was calculated as follows:  $[(a-A)/A] \times 100$ , Where: a: mean weight of treated larvae. A: mean weight of control larvae.
- **Pupation rate:** It was calculated according to Jimenez-Peydro *et al.* (1995) [27] as follows:  $PR = [\text{No. pupated larvae} / \text{No. treated larvae}] \times 100$
- **Developmental duration and rate:** The developmental durations (in days) were estimated using Dempster's equation (1957) [28]. The developmental rate was calculated using the Richard's equation (1957) [29].
- **Adult emergence:** The number of metamorphosed adults was expressed in % according to Jimenez-Peydro *et al.* (1995) [27] as follows:  $[\text{No. of completely emerged adults} / \text{No. of pupae}] \times 100$
- **Morphogenic efficiency:** It was calculated according to Vargas and Sehnal (1973) [30] as follows:  $[\text{No. of deformed adults} / \text{No. of emerged adults}] \times 100$  Adult longevity was determined in days, from the emergence until death.

### 4. Statistical analysis of data

Data obtained were analyzed by the Student's *t*-distribution, and refined by Bessel correction (Moroney, 1956) [31] for the test significance of difference between means  $\pm$  SD.

### Results

As clearly shown in the data of Table 1, Spinetoram considerably shortened the larval duration of the red palm weevil *Rhynchophorus ferrugineus* larvae, irrespective of the concentration. Larval duration was shortened to  $7.77 \pm 2.2$ ,  $8.41 \pm 2.0$ ,  $9.12 \pm 2.4$ , at 100, 50, 25 and 12.5 ppm, compared

to  $11.7 \pm 1.6$  days of control larvae. Only the highest concentration (4000 ppm) of Pyriproxyfen significantly shortened the larval duration in a developmental rate of 13.89, in comparison of 8.54% for controls. Moreover, Neemazal had no effect on the larval duration or developmental rate.

With regard to the larval body weight, it generally decreased, regardless the tested compound. Amongst all tested compounds, the strongest reducing effect on larval body weight was exhibited by Pyriproxyfen, at its higher concentration levels ( $1.98 \pm 0.63$ ,  $2.00 \pm 0.49$  and  $2.41 \pm 0.38$  mg, at 4000, 2000 and 1000 ppm, respectively, vs.  $3.89 \pm 0.74$  mg of control larvae with growth inhibition 49.1%.

Data of pupation and pupal duration had been arranged in Table (2). Depending on these data, pupation % decreased after treatment with all compounds. Treatment with the highest concentration led to reduction of pupation to 40, 50 and 10%, by Pyriproxyfen, Neemazal and Spinetoram, respectively. The pupal duration was shortened at all concentrations of the tested compounds, with exception of lower concentrations of Neemazal. As for example, the pupal duration was considerably shortened by Spinetoram to  $8.46 \pm 1.09$ ,  $12.14 \pm 3.01$  and  $12.33 \pm 3.66$  days, at 50, 25 and 12.5 ppm, compared to  $18.21 \pm 1.62$  days of control pupae. The developmental rate increased to 11.82, 8.24 and 8.11, at these concentration levels, compared to 5.49% of controls.

Table (3) contains results of the most important parameters of the adult performance. The adult emergence was detrimentally inhibited by Spinetoram (10.00, 50.00, 71.40 and 88.90% of adult emergence, at 100, 50, 25, and 12.5 ppm, respectively, vs. 90.00% of control adults). Also, Pyriproxyfen prohibited the adult emergence at all concentrations and the highest reduction was determined as 50.00% at 4000 ppm. On the other hand, Neemazal reduced the adult emergence at all concentrations, except at the lower two levels.

All insecticidal compounds caused some adult deformities. Fifteen percent of adult deformation had been caused by pyriproxyfen (at 4000, 2000, and 1000 ppm) and neemazal (at 40000 ppm). Also, spinetoram exhibited adult deformation, to some extent, at 50 and 12.5 ppm. However, the deteriorated adult morphogenesis was observed in atrophied wings and legs as well as expanded wings (see Fig. 1).

The tested compounds exhibited some shortening effects on the adult longevity. Adult longevity was remarkably shortened to  $41.62 \pm 3.56$  and  $44.67 \pm 5.86$  days, at 2000, 1000 ppm, respectively, of Pyriproxyfen; to  $55.34 \pm 7.81$  days, at 20000 ppm of neemazal;  $54.79 \pm 6.98$  days, at 25 ppm of spinetoram, compared to  $62.45 \pm 5.67$  days of control adults.

**Table 1:** Larval growth and development of *Rh. ferrugineus* after treatment of early 5<sup>th</sup> instar larvae with certain insecticidal compounds.

Compound	Conc. (ppm)	Duration (mean days $\pm$ SD)	Develop. Rate (%)	Weight (mean mg $\pm$ SD)	Growth inhibition (%)
Pyriproxyfen	4000	$7.2 \pm 2.97$ c	13.89	$1.98 \pm 0.63$ d	49.10
	2000	$10.8 \pm 1.03$ a	9.26	$2.00 \pm 0.49$ d	48.58
	1000	$11.0 \pm 1.16$ a	9.09	$2.41 \pm 0.38$ d	38.04
	500	$11.9 \pm 0.73$ a	8.40	$3.44 \pm 0.42$ a	11.58
	250	$11.3 \pm 0.94$ a	8.85	$3.54 \pm 0.47$ a	8.99
Neemazal	40000	$9.9 \pm 1.37$ a	10.10	$2.53 \pm 0.49$ c	34.96
	20000	$10.22 \pm 1.71$ a	9.78	$2.91 \pm 0.51$ b	25.19
	10000	$11.3 \pm 1.33$ a	8.85	$2.77 \pm 0.37$ c	28.79
	5000	$11.5 \pm 1.34$ a	8.70	$2.97 \pm 0.46$ c	23.65
	2500	$11.2 \pm 0.66$ a	8.93	$3.53 \pm 0.42$ a	9.25
Spinetoram	100	5*	25.77	2.36	39.33

	50	7.77 ± 2.2 c	17.33	2.5 ± 0.75 c	35.73
	25	8.41 ± 2.0 c	12.87	2.89 ± 0.81 b	25.70
	12.5	9.12 ± 2.4 b	10.96	2.86 ± 0.49 c	26.47
Control		11.7 ± 1.6	8.54	3.89 ± 0.74	---

**Conc:** Concentration; mean±SD followed with the letter (a): is not significantly different (P>0.05), (b): significantly different (P<0.05), (c): highly significantly different (P<0.01), (d): very highly significantly different (P<0.001). One insect.

**Table 2:** Pupation and developmental duration of pupae of *Rh. ferrugineus* after treatment of early 5<sup>th</sup> instar larvae with certain insecticidal compounds.

Compound	Conc. (ppm)	Pupation (%)	Pupal duration (mean days ± SD)	Develop. rate
Pyriproxyfen	4000	40	10.09 ± 6.26 c	9.91
	2000	60	13.0 ± 3.74 c	7.69
	1000	70	14.37 ± 2.50 c	6.96
	500	100	14.61 ± 2.41 c	6.84
	250	90	15.4 ± 1.64 c	6.49
Neemazal	40000	50	11.0 ± 6.04 c	9.09
	20000	50	11.5 ± 4.32 c	8.7
	10000	70	16.0 ± 4.32 a	6.25
	5000	90	18.77 ± 3.52 a	5.33
	2500	100	18.0 ± 1.48 a	5.56
Spinetoram	100	10	---	---
	50	40	8.46 ± 1.09 d	11.82
	25	70	12.14 ± 3.01 d	8.24
	12.5	90	12.33 ± 3.66 d	8.11
<b>Control</b>		100	18.21 ± 1.62	5.49

Conc., a, c, d: see footnote of Table 1.

**Table 3:** Adult performance of *Rh. ferrugineus* after treatment of early 5<sup>th</sup> instar larvae with certain insecticidal compounds.

Compound	Conc. (ppm)	Pupation (%)	Pupal duration (mean days±SD)	Develop. rate
Pyriproxyfen	4000	50.00	50.00	---
	2000	66.66	50.00	41.62 ± 3.56 d
	1000	57.71	50.00	44.67 ± 5.86 d
	500	40.00	14.28	58.44 ± 4.33 a
	250	88.88	---	64.51 ± 5.58 a
Neemazal	40000	40.00	50.00	---
	20000	80.00	25.00	55.34 ± 7.81 b
	10000	71.42	40.00	62.56 ± 5.66 a
	5000	90.00	28.50	61.78 ± 5.87 a
	2500	90.00	11.11	60.56 ± 5.89 a
Spinetoram	100	10.00	---	---
	50	50.00	25.00	55.62*
	25	71.40	---	54.79 ± 6.98 b
	12.5	88.90	12.50	59.52 ± 5.66 a
<b>Control</b>		90.00	---	62.45 ± 5.67

Conc., a, c, d: see footnote of Table 1.



**Fig 1:** The most important malformations of *Rh. ferrugineus* adults after treatment the early 5<sup>th</sup> instar larvae with certain insecticidal compounds. a): normal adult, b): pupal-adult intermediate, c): atrophied wings and legs, d): expanded wings.

**Discussion**

**1. Reduced body weight and growth**

The larval body weight of the red palm weevil *Rhynchophorus ferrugineus*, in the current investigation, was significantly reduced after treatment with all insecticidal compounds (pyriproxyfen, neemazal and spinetoram).

Similar results had been reported for prepupae and pupae of the same weevil by the plants products, Azadirachtin, jojoba oil, and the insect growth regulators (IGRs), Lufenuron and Diofenolan (Tanani, 2001) [32]. The present result was in accordance to some reducing effects different extracts and compounds on the body weight in different insects, such as the Egyptian cotton leafworm *Spodoptera littoralis* after treatment with Neemazal (Ghoneim *et al.*, 2000) [18], the false stable fly *Muscina stabulans* after treatment with Margosan-O (Al-Dali *et al.*, 2003) [33] and the house fly *Musca domestica* after treatment with Margosan-O (Amer *et al.*, 2004) [34], and the desert locust *Schistocerca gregaria* after treatment with neemazal (Hamadah, 2009) [35]. Reduction of the body weight, or weight gain, or the inhibition of growth, in general, by some IGRs, had been reported for several insect species, such as *M. domestica* (Bakr *et al.*, 1991) [36], the African armyworm *Spodoptera exempta*, the beet armyworm *Spodoptera exigua*, the cabbage moth *Mamestra brassicae* and the greater wax moth *Galleria mellonella* (Smaghe and Degheele, 1994) [37], *S. littoralis* (Ghoneim *et*

al., 1998)<sup>[38]</sup>, and the house mosquito *Culex pipiens* (Djeghader *et al.*, 2013, Djeghader *et al.*, 2014)<sup>[39, 40]</sup> by Novaluron, *C. pipiens* by Kinoprene (Hamaidia and Soltani, 2014)<sup>[41]</sup> and the black cutworm *Agrotis ipsilon* by Methoprene and Flufenoxuron (Khatter, 2014)<sup>[42]</sup>, as well as, *P. unionalis* by Novaluron (Ghoneim *et al.*, 2017)<sup>[43]</sup>. However, such suppressing action of the tested insecticidal compounds, in the present study, might be due to an ecdysonergic activity, as suggested by Smagghe and Degheele (1994)<sup>[37]</sup> after using tebufenozide (an ecdysone agonist).

## 2. Affected developmental durations

In the current study on *Rh. ferrugineus*, duration of the treated larvae was considerably shortened by all concentrations of Spinetoram but at the highest concentration of pyriproxyfen. To some extent, this result agreed with some reported results for other plant extracts and IGRs, such as *M. domestica* by Margosan-O (Ghoneim *et al.*, 2004)<sup>[44]</sup>, *S. gregaria* by Neemazal (Hamadah, 2009)<sup>[35]</sup>, *S. littoralis* (Ghoneim *et al.*, 2015)<sup>[45]</sup> and olive leaf moth *Palpita unionalis* (Ghoneim *et al.*, 2017)<sup>[43]</sup> by novaluron, and *M. stabulans* by pyriproxyfen (Hamadah, 2018)<sup>[46]</sup>. On the other hand, the pupal duration of *Rh. ferrugineus* was adversely affected by the tested insecticidal compounds in the present investigation. However, similar results on the same weevil had been reported for different compounds, such as neem and Flufenoxuron (El-Bokl *et al.*, 2010)<sup>[47]</sup>, lufenuron and diofenolan (Tanani, 2001)<sup>[32]</sup>, as well as the affected pupal duration of *M. stabulans* by pyriproxyfen (Hamadah, 2018)<sup>[46]</sup>. In contrast to our result, some authors reported prolongations in the pupal period in different insects by some IGRs, such as the vinegar fly *Drosophila melanogaster* by Pyriproxyfen (Bensebaa *et al.*, 2015)<sup>[48]</sup>, the mosquitoes *Culex quinquefasciatus* and *Aedes albopictus* after larval treatments with pyriproxyfen or methoprene (Khan *et al.*, 2016)<sup>[49]</sup> and the pink bollworm *Pectinophora gossypiella* after treatment of newly hatched or full grown larvae with Novaluron (Hassan *et al.*, 2017)<sup>[50]</sup>. To understand the shortening or prolongation of the developmental periods, it may be ascribed to the effects of IGRs on the release of ecdysteroids indirectly, by interfering with the neuroendocrine sites responsible for the release of tropic hormones (especially the prothoracicotrophic hormone) (Schluter *et al.*, 1985; Subrahmanyam *et al.*, 1989)<sup>[51, 52]</sup>.

## 3. Inhibited pupation and adult emergence

In the present study on *Rh. ferrugineus*, both pupation and adult emergence were significantly inhibited by the tested compounds. This result was, to a great extent, corroborated with those reported results for various compounds against different insects, since the adult emergence of *D. melanogaster* was inhibited by pyriproxyfen (Bensebaa *et al.*, 2015)<sup>[48]</sup>, both pupation and adult emergence percentages were reduced in the stable fly *Stomoxys calcitrans* by pyriproxyfen (Liu *et al.*, 2012)<sup>[53]</sup>. On the other hand, Tanani (2001)<sup>[32]</sup> recorded no effect on pupation% of RPW by Azadirachtin and jojoba oil. Also, pyriproxyfen suppressed the adult emergence in Asian citrus psyllid (Boina *et al.*, 2009)<sup>[54]</sup>, the mosquito *Aedes aegypti* (Sihuincha *et al.*, 2005; Harburger *et al.*, 2011)<sup>[55, 56]</sup>, the mosquitoes *Anopheles subpictus*, *Anopheles culicifacies* (Yapabandara and Curtis, 2004)<sup>[57]</sup>, the mosquito *Anopheles gambiae* (Mbare *et al.*, 2013)<sup>[58]</sup>, dark-winged fungus gnat *Bradysia coprophila*

(Ludwig and Oetting, 2001)<sup>[59]</sup>, the fly *Lycoriella ingenua* (Erler *et al.*, 2011)<sup>[60]</sup> and *Cx. pipiens* (Al-Sarar *et al.*, 2011)<sup>[61]</sup>. In addition, Singh and Kumar (2015)<sup>[62]</sup> reported an inhibition of adult emergence in the flesh fly *Sarcophaga ruficornis* by pyriproxyfen. Pupal treatment of the parasitic wasp *Encarsia formosa* with pyriproxyfen resulted in blockage of adult emergence (Wang and Liu, 2016)<sup>[63]</sup>. Adult emergence of *P. unionalis* was blocked by methoxyfenozide (Hamadah *et al.*, 2017)<sup>[64]</sup>. Moreover, adult emergence was completely blocked in the rice meal moth *Corcyra cephalonica* after treatment of 4<sup>th</sup> instar larvae with fenoxycarb (Singh and Tiwari, 2016)<sup>[65]</sup>. In general, inhibition of pupation and blockage of adult emergence might be result by the reduction of eclosion hormone production or release, since this hormone is responsible for some prerequisite process of the completion of moulting in insects (Ghoneim *et al.*, 1998)<sup>[38]</sup>.

## 4. Influenced adult longevity

The adult longevity of *Rh. ferrugineus* was shortened in the current study, irrespective of the tested compounds. Our result was in corroboration with those reported shortening effects on the adult longevity in different insects by various botanicals and IGRs, such as *S. littoralis* by the methanolic extract of *M. azedarach* (Schmidt *et al.*, 1997)<sup>[66]</sup>; *M. stabulans* by Margosan-O (Ghoneim and Al-Dali, 2002)<sup>[67]</sup>; *M. domestica* by lufenuron or diofenolan (Ghoneim *et al.*, 2004)<sup>[44]</sup> or Margosan-O (Amer *et al.*, 2004)<sup>[34]</sup>; *A. ipsilon* by pyriproxyfen, IKI-7899 or flufenoxuron (Shaurub *et al.*, 1999; El-Shiekh, 2002)<sup>[68, 69]</sup> and *M. stabulans* by Pyriproxyfen (Hamadah, 2018)<sup>[46]</sup>. On the contrary, El-Sayed (1982)<sup>[70]</sup> observed no effect of a neem seed suspension on the adult longevity of *S. littoralis*; and Hassan (2002)<sup>[71]</sup> recorded only a weak effect of the neem *Melia azedarach* extracts on the adult longevity of the same lepidopteran. The shortened adult longevity can be explained by the accumulation of toxic xenobiotics in the insect body which disturbs a complicated balance of factors such as absorption, excretion and detoxification (Abdel-Al, 1996)<sup>[72]</sup>.

## 5. Deteriorated adult morphogenesis

In the present study, disrupted morphogenesis was observed only in adults of *Rh. ferrugineus*, since some deformed adults had been recorded, irrespective of the tested compound. To some extent, the present finding was in partial resemblance with the reported results for various insect pests after using different compounds, such as the tsetse fly *Glossina morsitans* (Jordan *et al.*, 1979)<sup>[73]</sup>, *C. pipiens* (Bakr *et al.*, 1989)<sup>[74]</sup>, and *M. stabulans* (Basiouny, 2000)<sup>[75]</sup> after treatment with diflubenzuron; the confused flour beetle *Tribolium confusum* (El-Sayed *et al.*, 1984)<sup>[76]</sup>, *M. domestica* (Miller and Schmidtman, 1985)<sup>[77]</sup>, *M. stabulans* (Ghoneim *et al.*, 1992)<sup>[78]</sup> after treatment with Bay SIR-8514; *P. argyrostoma* (Ghoneim and Ismail, 1995)<sup>[79]</sup> and *M. stabulans* (Basiouny, 2000)<sup>[75]</sup> after treatment with IKI-7899 as well as *Rh. ferrugineus* after treatment with Azadirachtin, jojoba oil, lufenuron or diofenolan (Tanani, 2001)<sup>[32]</sup>; and *M. domestica* after treatment with lufenuron or diofenolan (Ghoneim *et al.*, 2004)<sup>[44]</sup> or with Jojoba (Hamadah, 2003)<sup>[80]</sup>. Recently, Hamadah (2018)<sup>[46]</sup> recorded some adult deformities of *M. stabulans* after treatment with pyriproxyfen. The adult deformation might be owing to the absence of necessary titer of ecdysteroids, as a response to

the action of botanicals and/or IGRs (Jagannadh and Nair, 1992) [81].

## Conclusion

In conclusion, the recorded disturbing effects of the tested insecticide alternatives (pyriproxyfen, neemazal, and spinetoram), in the present study, on growth, development, and adult performance of the red palm weevil, *Rhynchophorus ferrugineus* revealed that the bioinsecticide spinetoram was the most potent for controlling this dangerous weevil.

## 5. References

- Saleh MRA. Red palm weevil, *Rhynchophorus ferrugineus* Oliver is first recorded in Egypt and introduced in African continent. List No. 10634 Africa:Collection No.22563. International Institute of Entomology, London, 1992.
- Cox ML. Red palm weevil, *Rhynchophorus ferrugineus*, in Egypt. FAO, Plant Protection Bulletin. 1993; 41(1):30-31.
- Helal H, El-Sebay Y. Contribution on the wood borers attacking date palm trees and its control in Egypt. Egyptian Journal of Horticulture. 1994; 21(1):25-46.
- El-Sebay Y. Control of red palm weevil *Rhynchophorus ferrugineus* (Olivier) in Egypt. Egyptian Journal of Agricultural Research. 2004; 82(4):1581-1589.
- Merghem A. Susceptibility of the red palm weevil, *Rhynchophorus ferrugineus* (Olivier) to the green muscardine fungus, *Metarhizium anisopliae* (Metsch.) in the laboratory and in palm trees orchards. The Egyptian Journal of Biological Pest Control. 2011; 21(2):179-183.
- Merghem A, Bibars E. Spatial distribution of date palm trees borers with interest to red palm weevil, *Rhynchophorus ferrugineus* infestation levels at Ismailia governorate in Egypt. Egyptian Journal of Applied Science. 2014; 29(6):170-176.
- EPPO (European, Mediterranean Plant Protection Organization). Data sheets on quarantine pests-*Rhynchophorus ferrugineus*. EPPO Bulletin. 2008; 38:55-59.
- Abo El-Saad M, Ajlan A, Shawir M, Abdulsalam K, Rezk M. Comparative toxicity of four pyrethroid insecticides against red palm weevil, *Rhynchophorus ferrugineus* (Olivier) under laboratory conditions. Journal of Pest Control and Environmental Sciences. 2001; 9(3):63-76.
- Ranasinghe C, Fernando W, Zaneer S, Mubarak A. Analysis of residue levels in Kernel and nut water of coconut palms injected with Monocrotophos. Journal of the National Science Foundation of Sri Lanka. 2003; 31(3&4):431-436.
- Gloria S, Monroy O, Gómez J, Olgún E. Assessment of the hyper accumulating lead capacity of *Salvinia minima* using bioadsorption and intracellular accumulation factors. Water, Air and Soil Pollution. 2008; 194(1-4):77-90.
- Koul O, Walia S. Comparing impacts of plant extracts and pure allelochemicals and implications for pest control. Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources, CAB Reviews. 2009; No. 049 (<http://www.cabi.org/cabreviews>).
- Smet H, Rans M, Loof AD. Comparative effectiveness of insect growth regulators with juvenile hormone, anti-juvenile hormone and chitin synthesis inhibiting activity against several stored food insect pests. In: (Fleurat-Lessard, F., Ducon, P., eds.). Proceedings of the 5th International Working Conference on Stored Product Protection, France, 1990, 649-657.
- Oberlander H, Silhacek DL, Shaaya E, Ishaaya I. Current status and future prospective of the use of insect growth regulators for the control of stored product insects. Journal of Stored Products Research. 1997; 33(1):1-6.
- Rembold H, Sieber KP. Inhibition of oogenesis and ovarian ecdysteroid synthesis by azadirachtin in *Locusta migratoria migratorioides*. Zeitschrift für Naturforschung. 1981; 36:466-469.
- Garcia EDS, Rembold H. Effect of azadirachtin on ecdysis of *Rhodnius prolixus*. Journal of Insect Physiology. 1984; 30:939-941.
- Dorn A, Rademacher M, Sehn E. Effects of azadirachtin on the moulting cycle, endocrine system, and ovaries in last instar larvae of the milk weed bug *Oncopeltus fasciatus*. Journal of Insect Physiology. 1986; 32:231-238.
- Richer K, Bohm GA, Kleeberg H. Effect of Neemazal, a natural azadirachtin-containing preparation, on *Periplaneta americana* (L.) (Orthoptera: Blattellidae). Journal of Applied Entomology. 1997; 121:59-64.
- Ghoneim KS, Mohamed HA, Bream AS. Efficacy of the neem seed extract, Neemazal, on the growth and development of the Egyptian cotton leafworm, *Spodoptera littoralis* Bois. (Lepidoptera: Noctuidae). Journal of Egyptian German Society of Zoology. 2000; 33:161-179.
- Hamadah SH, Tanani MA. Laboratory studies to compare the toxicity for three insecticides in the red palm weevil *Rhynchophorus ferrugineus*. International Journal of Biology, Pharmacy and Allied Science. 2014; 2(3):506-519.
- Hamadah KH, Tanani MA. Disruptive Impacts of Selected Insecticides on Larval Haemogram parameters of the Red Palm Weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). Egyptian Academic Journal of biological Sciences. 2017; 10(8):85- 97.
- Sparks TC, Thompson GD, Kirst HA, Hertlein MB, Larson LL, Worden TV, Thibault ST. Biological activity of the spinosyns, new fermentation derived insect control agents, on tobacco budworm (Lepidoptera: Noctuidae) larvae. Journal of Economic Entomology. 1998; 91:1277-1283.
- Watson GB. Action of insecticidal spinosyns on  $\gamma$ -aminobutyric acid responses from small-diameter cockroach neurons. Pesticide Biochemistry and Physiology. 2000; 71:20-28.
- Sparks TC. New insect control agents: mode of action and selectivity, In: The management of diamondback moth and other crucifer pests (Endersby, N.M. and Ridland, P.M., eds.). Proceedings of the fourth international workshop, 26-29, Melbourne. Department of Natural Resources and Environment, Melbourne, Australia, 2004-2001, 37-44.
- Sarfraz M, Dossdall LM, Keddie BA. Spinosad: A promising tool for integrated pest management. Outlooks Pest Management. 2005; 16:78-84.
- Rahalkar GW, Harwalkar MR, Ranavavare HD. Development of red palm weevil, *Rhynchophorus*

- ferrugineus* Oliv. on sugarcane. Indian Journal of Entomology. 1972; 34:213-215.
26. Ranavare HD, Shantharan K, Harwalkar MR, Rahalhar GW. Method for laboratory rearing of red palm weevil, *Rhynchophorus ferrugineus* (Oliv.). Journal of Plant Crops. 1975; 3:65-67.
  27. Jimenez-Peydro R, Gimeno-Martos C, Lopez-Ferrer J, Serrano-Delgado C, Moreno-Mari J. Effects of the insect growth regulator, cyromazine, on the fecundity, fertility and offspring development of Mediterranean fruit fly, *Ceratitis capitata* Wied (Diptera, Tephritidae). Journal of Applied Entomology. 1995; 119:435-438.
  28. Dempster CV. The population dynamic of Moroccan locust *Dociostarus murcocamus* in Cyprus. AntiLocust Bulletin, 1957, 27.
  29. Richard AG. Cumulative effects of optimum and suboptimum temperatures on insect development. In: Influence of Temperature on Biological Systems (Johnson, F.H., ed.). American Physiological Society. 1957; 15:35-38.
  30. Vargas L, Sehna F. Use of juvenile hormone analogue against the fall webworm, *Hyphanthia cunea*. Entomologia Experimentalis et Applicata. 1973; 16:115-122.
  31. Moroney MJ. Facts from figures (3rd ed.). Penguin Books Ltd., Harmondsworth. Middle Sex, 1956.
  32. Tanani AM. Study the effects of certain IGRs and plant extracts on some physiological aspect of the *Rhynchophorus ferrugineus* (Curculionidae: Coleoptera). M.Sc. Thesis, Faculty of Science, Al-Azhar University, Cairo, Egypt, 2001.
  33. Al-Dali AG, AbdEl-Ghaffar A, Ghoneim KS. Efficacy of the neem seed extract, Margosan-O, on the development and morphogenesis of *Muscina stabulans* (Diptera: Muscidae). Journal of Egyptian German Society of Zoology. 2003; 42:85-100.
  34. Amer MS, Ghoneim KS, Al-Dali AG, Bream AS, Hamadah KhSh. Assessment of the activity of Margosan-O and Jojoba against the house fly *Musca domestica* (Diptera: Muscidae). Al-Azhar Bulletin of Science. 2004; 15(2):9-24.
  35. Hamadah KH. Some developmental, haematological and enzymatic effects of certain plant extracts on the desert locust *Schistocerca gregaria* (Orthoptera: Acrididae). Ph.D. Thesis, Faculty of Science, Al-Azhar University, Cairo, Egypt, 2009.
  36. Bakr RFA, Abd El-Razek NA, Hamed MS, Guneidy AM. Physiological effect of some insect growth regulators on the respirometric measurements, total protein and free amino acids of the house fly *Musca domestica*. Ain Shams Bulletin of Science. 1999; 28(B):169-183.
  37. Smagghe G, Degheele D. Action of a novel nonsteroidal ecdysteroid mimic, tebufenozide (RH-5992), on insects of different orders. Pesticide Science. 1994; 42(2):85-92.
  38. Ghoneim KS, Bream AS, Mohamed HA. Bioactivity of the ecdysteroid agonist Tebufenozide (RH-5992) on the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.) (Noctuidae: Lepidoptera). Al-Azhar Bulletin of Science. 1998; 9(2):947-963.
  39. Djeghader N, Djeghader H, Bouaziz, A, Soltani N. Biological effects of a benzoylphenylurea derivative (Novaluron) on larvae of *Culex pipiens* (Diptera: Culicidae). Advances in Applied Science Research. 2013; 4(4):449-456.
  40. Djeghader NEH, Aïssaoui L, Amira K, Boudjelida H. Impact of a chitin synthesis inhibitor, Novaluron, on the development and the reproductive performance of mosquito *Culex pipiens*. World Applied Science Journal. 2014 ; 29(7):954-960.
  41. Hamaidia K, Soltani N. Laboratory evaluation of a biorational insecticide, Kinoprene, against *Culex pipiens* larvae: effects on growth and development. Annual Research and Review in Biology. 2014; 4(14):2263-2273.
  42. Khatter NA. Effect of two insect growth regulators on the development of *Agrotis ipsilon* Hufn. (Lepidoptera: Noctuidae). Journal of Harmonized Research and Applied Science. 2014; 2(1):20-28.
  43. Ghoneim K, Hamadah Kh, Mansour A, Abo Elsoud A. Toxicity and disruptive impacts of Novaluron, a chitin synthesis inhibitor, on development and metamorphosis of the olive leaf moth *Palpita unionalis* (Hübner) (Lepidoptera: Pyralidae). International Journal of Trend in Research Development. 2017 ; 4(3):184-193.
  44. Ghoneim KS, Amer MS, Bream AS, Al-Dali AG, Hamadah Kh Sh. Developmental and morphogenic responses of the house fly *Musca domestica* to the CSIs: Lufenuron and Diufenolan. Al-Azhar Bulletin of Science. 2004; 15(2):25-42.
  45. Ghoneim K, Tanani MA, Hamadah Kh, Basiouny A, Waheeb H. Bioefficacy of Novaluron, a chitin synthesis inhibitor, on survival and development of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). Journal of Advanced Zoology. 2015; 1(1):24-35.
  46. Hamadah Kh Sh. Impaired survival and development of the false stable fly, *Muscina stabulans* (Fallen) (Diptera: Muscidae) by Pyriproxyfen (a juvenile hormone analogue). Journal of Egyptian Society Parasitology. 2018; 48(3):677-688.
  47. El-Bokl MM, Baker RFA, El-Gammal HL, Mahmoud MZ. Biological and histopathological effects of some insecticidal agents against red palm weevil *Rhynchophorus ferrugineus*. Egyptian Academic Journal of biological Sciences. 2010; 1(1):7-22.
  48. Bensebaa F, Kilani-Morakchi S, Aribi N, Solatani N. Evaluation of Pyriproxyfen, a juvenile hormone analog, on *Drosophila melanogaster* (Diptera: Drosophilidae): Insecticidal activity, ecdysteroid contents and cuticle formation. European Journal of Entomology. 2015; 112(4):625-631.
  49. Khan GZ, Khan I, Khan IA, Alamze b, Salman M, Ullah K. Evaluation of different formulations of IGRs against *Aedes albopictus* and *Culex quinquefasciatus* (Diptera: Culicidae). Asian Pacific. Journal of Tropical Biomedicine. 2016; 6(6):485-491.
  50. Hassan HA, Ghoneim K, Tanani MA, Bakr NA. Impairing effectiveness of the chitin synthesis inhibitor, Novaluron, on adult performance and reproductive potential of the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). Journal of Entomology and Zoology Studies. 2017; 5(2):581-592.
  51. Schluter U, Bidmon HJ, Grewe S. Azadirachtin affects growth and endocrine events in larvae of the tobacco hornworm *Manduca sexta*. Journal of Insect Physiology. 1985; 31:773-777.
  52. Subrahmanyam B, Muller T, Rembold H. Inhibition of turnover of neurosecretion by azadirachtin in *Locusta*

- migratoria*. Journal of Insect Physiology. 1989; 35:493-500.
53. Liu SS, Li AY, Lohmeyer KH, Perezdeleon AA. Effects of Pyriproxyfen and buprofezin on immature development and reproduction in the stable fly. Medical and Veterinary Entomology. 2012; 26(4):379-85.
  54. Boina D.R, Rogers ME, Wang N, Stelinski, L. Effect of Pyriproxyfen, a juvenile hormone mimic, on egg hatch, nymph development, adult emergence and reproduction of the Asian citrus psyllid, *Diaphorina citri* Kuwayama. Pest Management Science. 2009; 66:349–357.
  55. Sihuinchu M, Zamora-Perea E, Orellana-Rios W, Stancil JD, López-Sifuentes V, Vidal-Oré C Devine GJ. Potential use of Pyriproxyfen for control of *Aedes aegypti* (Diptera:Culicidae) in Iquitos, Perú. Journal of Medical Entomology. 2005; 42:620–630.
  56. Harburger L, Beltrán G, Goldberg L, Goldberg L, Zerba E, Licastro, S, Masuh H. A new strategy for *Aedes aegypti* (Diptera:Culicidae) control with community participation using a new fumigant formulation. Journal of Medical Entomology. 2011; 48:577–583.
  57. Yapabandara AMGM, Curtis CF. Control of vectors and incidence of malaria in an irrigated settlement scheme in Sri Lanka by using the insect growth regulator Pyriproxyfen. Journal of the American Mosquito Control Association. 2004; 20:395-400.
  58. Mbare O, Lindsay SW, Fillinger U. Dose-response tests and semi-field evaluation of lethal and sub-lethal effects of slow release Pyriproxyfen granules (Sumilarv®0.5G) for the control of the malaria vectors *Anopheles gambiae* sensu lato. Malaria Journal. 2013; 12:94.
  59. Ludwig SW, Oetting RD. Evaluation of medium treatments for management of *Frankliniella occidentalis* (Thripidae:Thysanoptera) and *Bradysia coprophila* (Diptera:Sciaridae). Pest Management Science. 2001; 57:1114–1118.
  60. Erler F, Polat E, Demir H, Catal M, Tuna G. Control of mushroom sciarid fly *Lycoriella ingenua* populations with insect growth regulators applied by soil drench. Journal of Economic Entomology. 2011; 104:839–844.
  61. Al-Sarar AS, Al-Shahrani D, Bayoumi AE, Abobakr Y, Hussein HI. Laboratory and field evaluation of some chemical and biological larvicides against *Culex* spp. (Diptera:Culicidae) immature stages. International Journal of Agriculture and Biology. 2011; 13:115–119.
  62. Singh S, Kumar K. Effects of juvenoid Pyriproxyfen on reproduction and F1 progeny in myiasis causing flesh fly *Sarcophaga ruficornis* L. (Sarcophagidae:Diptera). Parasitology Research. 2015; 114:2325–2331.
  63. Wang QL, Liu TX. Effects of three insect growth regulators on *Encarsia formosa* (Hymenoptera:Aphelinidae), an endoparasitoid of *Bemisia tabaci* (Hemiptera:Aleyrodidae). Journal of Economic Entomology. 2016 ; 1-8.
  64. Hamadah Kh, Ghoneim K, Mansour A, Abo Eloud A. Deranged adult performance and reproductive potential of the olive leaf moth *Palpita unionalis* (Hubner) (Lepidoptera:Pyralidae) by the non-steroidal ecdysone agonist, Methoxyfenozide. International Journal of Information Research and Review. 2017; 4(6):4228-4240.
  65. Singh A, Tiwari SK. Role of Fenoxycarb, a juvenile hormone analogue, on the developmental stages of rice-moth, *Corcyra cephalonica* Staint. (Lepidoptera:Pyralidae). International Journal of Zoological Investigations. 2016; 2(2):267-280.
  66. Schmidt GH, Ahmed AAI, Breuer M. Effect of *Melia azedarach* extract on larval development and reproduction parameters of *Spodoptera littoralis* (Boisd.) and *Agrotis ipsilon* (Hufn.) (Lepidoptera:Noctuidae). Anzeiger-Fur-Schadlingskunde, Pflanzenschutz, Umweltschutz. 1997; 70 (1):4-12.
  67. Ghoneim KS, Al-Dali AG. Survival and reproductive responses of *Muscina stabulans* Fallen (Muscidae:Diptera) to the neem seed extracts, Margosan-O. 2nd International Conference of Plant Protection Research Institute, Cairo, Egypt. 2002; 21- 24 December.
  68. Shaurub EH, Emara SA, Zohdy, NZ, Abdel-Aal AE. Effect of four insect growth regulators on the black cutworm. *Agrotis ipsilon* (Hufn.) (Lepidoptera: Noctuidae). The International Conference of Pest Control, Mansoura University, Egypt September. 1999; 733-776.
  69. El-Sheikh TAA. Effects of application of selected insect growth regulators and plant extracts on some physiological aspects of the black cutworm, *Agrotis ipsilon* (Huf.). Ph.D. Thesis, Faculty of Science, Ain Shams University, Cairo, Egypt. 2002; 189pp.
  70. El-Sayed EI. Neem *Azadirachta indica* A. Juess, seeds as antifeedant and ovipositional repellent for the Egyptian Cotton leafworm, *Spodoptera littoralis* (Boisd.). Bulletin of Entomological Society of Egypt (Econ. Ser.). 1982; 15:49-58.
  71. Hassan HA. Biological and biochemical studies on the effects of some botanical extracts on cotton leafworm *Spodoptera littoralis* (Boisd.) (Lepidoptera:Noctuidae). Unpublished M.Sc.Thesis, Faculty of Science, Ain Shams University, Cairo, Egypt. 2002.
  72. Abdel-Al AE. Biological, histological and physiological effects of some insect growth regulators on the greasy cutworm, *Agrotis ipsilon* (Lepidoptera:Noctuidae). M.Sc.Thesis, Faculty of Science,Cairo University Egypt. 1996.
  73. Jordan AM, Trewern MA, Borkovec AB, De Milo AB. Laboratory studies on the potential of three insect growth regulators for control of tsetse fly, *Glossina morsitans* Westwood (Dipt., Glossinidae). Bulletin of Entomological Research. 1979; 69:55-64.
  74. Bakr RFA, Abo Gabal NM, Hussein MA. Insect growth regulators:I. Biological activity of some IGRs against the susceptible and resistant strains of *Culex pipiens* larvae. Journal of Egyptian Society of Parasitology. 1989; 19(2):589-597.
  75. Basiouny A.L. Some physiological effects of certain insect growth regulators on the false stable fly, *Muscina stabulans* (Fallen.) (Diptera:Muscidae). Unpublished Ph.D. Thesis, Faculty of Science, Al-Azhar University, Cairo, Egypt. 2000.
  76. El-Sayed MA, Abdel-Razik F, Kandil MA. Biological activity of the insect growth regulator triflumuron against *Tribolium confusum*. Bulletin of Entomological Society of Egypt (Econ. Ser.). 1984;14:171-175.
  77. Miller RW, Schmidtman ET. Feeding cyromazin (Lerodex) to dairy calves for the control of house flies. Journal of Agricultural Entomology. 1985; 2:31-36.
  78. Ghoneim KS. Essa N, Abul-Ela RG, Al-Morsy AA,

- Nassar Ml. Efficacy of Triflumuron (Bay Sir-8514) for remedial control of the false stable fly *Muscina stabulans* (Fallen) (Diptera:Muscidae). Al-Azhar Bulletin of Science.1992; 3 (2):687 – 693.
79. Ghoneim KS, Ismail IE. Survival, developmental and morphogenic deficiencies of *Parasarcophaga argyrostoma* (Diptera: Sarcophagidae) induced by the chitin biosynthesis inhibitor, chlorfluazuron (IKI-7899). Journal of Egyptian Society of Parasitology. 1995; 25(2):561–581.
80. Hamadah KH. Physiological and Biochemical Effects of IGRs and plant extracts on the house fly *Musca domestica*. M.Sc. Thesis, Faculty of Science, Al-Azhar University, Cairo, Egypt, 2003.
81. Jagannadh V, Nair VSK. Azadirachtin-induced effects on larval-pupal transformation of *Spodoptera mauritia*. Physiological Entomology. 1992; 17:56-61.