



Inhibitory impact of Novaluron, a recent insect growth regulator, on the reproductive potential of *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae)

Basiouny A^{1*}, Waheeb H²

^{1,2} Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Madenit Nasr, Cairo, Egypt

Abstract

The Egyptian cotton leafworm *Spodoptera littoralis* is one of the destructive pests of field crops in the tropical and subtropical areas of the world. The objective of the present study was to assess the impact of Novaluron on the reproductive potential of this pest. A series of concentrations (1.0, 0.1, 0.01, 0.001 & 0.0001 ppm) was prepared and applied on the freshly molted penultimate instar larvae. Another series (0.1, 0.01, 0.001 & 0.0001 ppm) was prepared and applied on the newly molted last (6th) instar larvae. The results can be summarized as follows. The tested compound exhibited a predominantly inhibitory effect on the oviposition efficiency, regardless the concentration and time of larval treatment. Both fecundity and fertility had been drastically reduced, in a dose-dependent course, regardless the time of treatment. Irrespective of the concentration and time of treatment, Novaluron halted the embryonic development, since the incubation period of eggs was remarkably prolonged. In the light of the present study, Novaluron acts as an antigonadotropic compound in *S. littoralis*.

Keywords: antigonadotropic, embryonic development, fecundity, fertility, incubation, oviposition

Introduction

The Egyptian cotton leafworm *Spodoptera littoralis* Boisduval (Lepidoptera: Noctuidae) represents a destructive pest in different parts of the world (Hill, 1987) [1]. It attacks plants belonging to more than forty families of varying economic importance (Lal and Naji, 1990) [2]. As for examples, it damages the glasshouses crops and flower production in Southern Europe, various types of vegetables in North Africa and cotton in Egypt (El-Aswad *et al.*, 2003; Roques *et al.*, 2008) [3,4]. The intensive use of broad-spectrum pesticides against *S. littoralis* has led to different problems, such as the development of resistance to many pesticides making its control even more difficult (Smaghe *et al.*, 1999; Miles and Lysandrou, 2002; Aydin and Gurkan, 2006) [5,6,7]. This insect has been subjected to extensive research, much of which is envisioned to finding new efficient ways to control (Hussain, 2012) [8].

Over few decades, the insect growth regulators (IGRs) have been considered as a possible alternative of the conventional insecticides for controlling this dangerous insect (Raslan, 2002) [9]. IGRs have novel modes of action which disrupt the development and some of other physiological processes of the target pest. These compounds tend to be selective and less toxic to non-target organisms than the conventional pesticides (Gurr *et al.*, 1999) [10]. Depending on the mode of action, IGRs had been recently categorized in substances interfering with the functions of insect hormones and chitin synthesis inhibitors (CSIs) (Tunaz and Uygun, 2004) [11]. In insects, CSIs interfere with the chitin biosynthesis and thus prevent moulting, or produce deformed cuticle (Hammock and Quistad, 1981) [12]. CSIs had experimentally proved to act as effective inhibitors of the development of insect pests (Ishaaya *et al.*, 2002, 2003; Kandil *et al.*, 2012) [13,14,15]. CSIs also affect the hormonal balance in insects, thereby resulting in various physiological disturbances (Soltani *et al.*, 1984) [16].

As a relatively new benzoylphenyl urea CSI, Novaluron

exhibits a remarkable activity against several insects, such as *Leptinotarsa decemlineata* (Cutler *et al.*, 2007; Alyokhin *et al.*, 2009) [17, 18], *Bemisia tabaci* and *Trialeurodes vaporariorum* (Kim *et al.*, 2000) [19] and *Helicoverpa* species (Ishaaya *et al.*, 2001, 2002, 2003) [20, 13, 14]. According to many authors (Barazani, 2001; Ishaaya and Horowitz, 2002; Ishaaya *et al.*, 2001, 2002) [21, 22, 20, 13], this CSI has probably a mild effect on the natural enemies and has no serious effect on parasitoids as well as has low mammalian toxicity. As recorded in Egypt by Malhata *et al.* (2014) [23], Novaluron residues tend to dissipate with a half-life of 2.08 days and the safe use of it on tomato, and probably on other crops. The objective of the current study was to evaluate the disruptive effects of Novaluron on the most important reproductive criteria of *S. littoralis*.

Materials and Methods

1. The insect under study

A culture of sensitive strain of *Spodoptera littoralis* (Lepidoptera: Noctuidae) was established at Faculty of Science, Al-Azhar University, Egypt, under controlled conditions (27±2°C, 65±5% R.H., photoperiod 14 h L and 10 h D). This culture was originated by a sample of pupae kindly obtained from susceptible strain maintained for some generations in Plant Protection Research Institute, Doqqi, and Giza, Egypt. The rearing procedure was achieved according to Ghoneim (1985) [24] and improved according to Bakr *et al.* (2010) [25]. Every day, larvae had been provided with fresh castor bean leaves *Ricinus communis*. The emerged adults were supplemented with 10% honey solution on a cotton wick as a food source. Moths were allowed to lay eggs on *Oleander* branches, and the egg patches were collected daily, and transferred into Petri dishes for another generation.

2. Novaluron and larval treatments

The tested CSI, Novaluron (1-[3-chloro-4-(1, 1, 2-tri fluoro-2-tri fluoro- methoxy- ethoxy) phenyl]-3-(2, 6-di fluorobenzoyl)

urea) (Rimon, Chemtura Corporation, Middlebury, CT), was supplied by Sigma-Aldrich Chemicals (<https://www.sigmaaldrich.com>). A series of concentrations was prepared using distilled water. Freshly moulted 5th (penultimate) instar larvae were treated with the concentrations 1.0, 0.1, 0.01, 0.001 & 0.0001 ppm, but the newly moulted 6th (last) instar larvae were treated with the concentrations 0.1, 0.01, 0.001 & 0.0001 ppm. The feeding larvae had been provided with fresh castor bean leaf discs after dipping in each concentration for 5 minutes. They were left to feed on the treated leaf discs for 24 hrs and then on untreated fresh leaf discs every day. Control larvae of the same age were allowed to feed on water-treated leaf discs. Ten replicates of treated and control larvae (one larva/replicate) were kept separately in glass vials. All larvae (control and treated) were carefully handled until the adult eclosion just after which all reproductive data were recorded.

3. Reproductive criteria

After 24 h of emergence, the treated adult females were kept separately in glass jars (1 L) and coupled with healthy adult males (1:2) of the same age obtained from the normal culture. Each jar was provided with sterilized cotton pieces soaked in 10% honey solution for feeding and fresh *Nerium oleander* branches as oviposition sites. During the oviposition period, egg-patches were collected daily and carefully transferred to Petri dishes to count eggs. Oviposition rate was calculated as follows: Number of laid eggs per ♀/reproductive lifetime (in days). Eggs were counted for calculating the number of eggs per female (Fecundity). The laid eggs were kept in Petri dishes under the same controlled conditions, as previously mentioned. Eggs were observed until hatching to calculate the incubation period (in days). The hatchability (Fertility) was usually expressed in hatching percentage of eggs. Sterility index was calculated according to Topozada *et al.* (1966)^[26] as follows: Sterility Index = $100 - [(a \ b / A \ B) \times 100]$. Where: a: mean number of eggs laid per female in the treatment. b: percentage of hatching in the treatment. A: mean number of eggs laid per female in the controls. B: percentage of hatching in the controls.

4. Statistical analysis of data

The present results were analyzed using the Student's *t*-distribution, and refined by Bessel correction (Moroney, 1956)^[27] for the test significance of difference between means.

Results

1. Effect of Novaluron on the oviposition efficiency of *S. littoralis*

In the present study, penultimate (5th) or last (6th) instar larvae of *S. littoralis* were treated with Novaluron at a dose range 1.0-0.0001 ppm. Because the oviposition rate in insects can be used as an informative indicator of the oviposition efficiency, Data arranged in Table (1) clearly reveal a

predominant inhibitory effect of Novaluron on the oviposition efficiency, regardless the concentration applied on 5th instar larvae. However, the severely declined oviposition rate was determined at the higher two concentrations (24.63±1.37 and 16.14±2.04, respectively, vs. 192.64±4.33 of control females). A similar suppressing action of Novaluron on the oviposition rate was recorded after treatment of 6th instar larvae, regardless the concentration, as obviously demonstrated in Table (2). Depending on this table, the strongest inhibitory effect of Novaluron was exhibited at the highest concentration (108.33±4.17 vs. 192.64±4.33 of control adult females).

2. Disrupted reproductive capacity of *S. littoralis* by Novaluron

In the light of data assorted in Table (1), treatment of 5th instar larvae of *S. littoralis* with Novaluron resulted in seriously inhibited fecundity, in a dose-dependent manner. With regard to the higher two concentrations, fecundity dropped (78.00±22.14 and 100.00±11.00 eggs/treated ♀, respectively, vs. 1733.40±57.23 eggs/control ♀). Also, strongly reducing action of Novaluron on the female fecundity was exerted after treatment of last instar larvae, in a dose-dependent course. The most reducing action was exerted at the highest concentration (787.50±109.60 eggs/treated ♀ vs. 1733.40±57.23 eggs/control ♀, table 2).

Data contained in Table (1) unexceptionally reveal a deteriorating action of Novaluron on fertility (egg viability) after treatment of 5th instar larvae. Although the reduction of fertility could not be detected in certain trend, the most potent reducing action of Novaluron was exerted at 1.00 and 0.01 ppm (37.23±0.00 and 44.70±4.24%, respectively, in comparison with 97.80±0.78% hatching eggs of controls). Moreover, the sterility index was found in a dose-dependent course. A similar fertility reduction was, also, determined after treatment of 6th instar larvae with Novaluron. In addition, the highest sterility index was calculated in 86.90 at the highest concentration (for detail, see table 2).

3. Disturbed embryonic development of *S. littoralis* by Novaluron

The incubation period of insect eggs can be used as a good indicator of the embryonic developmental rate, i.e., shorter period usually denotes faster developmental rate and *vice versa*. After treatment of 5th instar larvae with Novaluron, data of the incubation period were distributed in Table (1). The embryonic development in *S. littoralis* had been generally subjected to a prolonged inhibitory effect of the tested compound since the incubation period was remarkably prolonged. At the highest concentration, the longest period was estimated (4.00±0.00 days, compared to 2.20±0.45 days of control eggs). Also, similar retarding action of Novaluron on the embryonic development was exerted after treatment of last instar larvae since the incubation period was significantly lengthened, regardless the concentration (see Table 2).

Table 1: Oviposition efficiency and reproductive potential of *S. littoralis* as affected by the treatment of newly moulted penultimate instar larvae with Novaluron

Conc. (ppm)	Oviposition rate	Fecundity (mean eggs±SD)	Fertility (%)	Sterility Index	Incubation period (mean days±SD)
1.00	024.63±1.37 d	078.00±22.14 d	37.23±0.00 d	98.33	4.00±0.00 d
0.10	016.14±2.04 d	100.00±11.00 d	72.00±0.00 d	95.83	4.00±0.00 d
0.01	140.08±4.12 d	721.00±18.39 d	44.70±4.24 d	81.00	4.00±0.00 d
0.001	132.58±3.67 d	945.00±76.88 d	65.60±0.57 d	63.45	3.00±1.00 d

0.0001	168.57±3.25 c	1128.85±62.94 d	84.20±1.94 d	43.91	3.43±0.53 d
Control	192.64±4.33	1733.40±57.23	97.80±0.78	---	2.20±0.45

Conc.: concentration level. Mean ± SD followed with the same letter (a): insignificantly different ($P > 0.01$), (b): significantly different ($P < 0.05$), (c): highly significantly different ($P < 0.01$), (d): very highly significantly different ($P < 0.001$).

Table 2: Oviposition efficiency and reproductive potential of *S. littoralis* as affected by the treatment of newly moulted last instar larvae with Novaluron.

Conc. (ppm)	Oviposition rate	Fecundity (mean eggs ± SD)	Fertility (%)	Sterility Index	Incubation period (mean days ± SD)
0.10	108.33±4.17 d	787.50±109.60 d	66.85±1.91 d	86.90	4.00±0.00 d
0.01	150.79±5.67 c	1251.67±76.54 d	75.43±17.67 d	44.38	3.67±0.58 d
0.001	173.39±4.51 b	1378.75±124.19 d	84.88±2.18 d	30.90	3.25±0.50 d
0.0001	152.81±3.97 c	1149.17±135.63 d	82.53±1.14 d	44.13	3.33±0.52 d
Control	192.64±4.33	1733.40±57.23	97.80±0.78	---	2.20±0.45

Conc., a, b, c, d: See footnote of Table (1)

Discussion

The insect growth regulators (IGRs)-treated larvae may develop as deformed adults who would be non-viable or with the least reproductive capacity (Williams and Amos 1974)^[28]. This result may be due to sterility or reduced fecundity in the treated insects (Ghoneim *et al.*, 2014)^[29]. However, disruptive impacts of IGRs on the reproduction in insects can be categorized into: i) deteriorated reproductive behaviour, ii) inhibited oviposition, iii) reduced hatchability of eggs, and iv) adult sterilization (Mondal and Parween, 2000)^[30].

1. Inhibited oviposition efficiency of *S. littoralis* by Novaluron

In insects, the oviposition efficiency can be indicated by the oviposition rate. In the current study, novaluron exerted a predominant inhibitory action on the oviposition efficiency of *S. littoralis*, irrespective of the time of larval treatment or concentration. This result was coincided with the reported inhibition of oviposition efficiency of the pink bollworm *Pectinophora gossypiella* after treatment of newly hatched and full grown larvae with novaluron (Hassan *et al.*, 2017)^[31] as well as reported results for other insects, such as *S. littoralis* by Tebufenozide (Bakr *et al.*, 2005)^[32], and flufenoxuron (Bakr *et al.*, 2010)^[25]; the desert locust *Schistocerca gregaria* by flufenoxuron and lufenuron (Soltani-Mazouni and Soltani, 1994)^[33] or tebufenozide (Al-Dali *et al.*, 2008)^[34]; the Indianmeal moth *Plodia interpunctella* by the ecdysteroid agonist RH-5849 (Smaghe and Degheele, 1994)^[35] and the cowpea weevil *Callosobruchas maculatus* by cyromazine (Al-Mekhlafi *et al.*, 2011)^[36]. This result was also in agreement with those reported results of inhibited oviposition of the olive leaf moth *Palpita unionalis* after treatment of newly moulted last instar larvae with methoxyfenozide (Hamadah *et al.*, 2017)^[37] and *P. gossypiella* after treatment of 1-day old eggs with Noviflumuron or Novaluron (Tanani and Ghoneim, 2018)^[38]. In contrast, the current result disagreed with the enhanced oviposition of the field cricket *Gryllus bimaculatus* by some ecdysteroids (Behrens and Hoffmann, 1983)^[39]. The suppressed oviposition, in the current study, may be understood by the inhibition of ovarian DNA synthesis or interference of the tested compound with vitellogenesis via certain biochemical processes. However, this IGR might exert a reverse action to those exhibited by the ecdysteroids which induce the neurosecretory cells to release a myotropic ovulation hormone (Smaghe *et al.*, 1996; Parween *et al.*, 2001)^[40, 41].

2. Perturbation of the reproductive capacity of *S. littoralis* by Novaluron

Reduced fecundity: There are many reported results of reduced fecundity of *S. littoralis* after treatment of larvae with various IGRs, such as diflubenzuron (Aref *et al.*, 2010)^[42], methoxyfenozide (Pineda *et al.*, 2009)^[43] and lufenuron (Abdel-Rahman *et al.*, 2007; Gaaboub *et al.*, 2012)^[44, 45]. Also, fecundity of other insects was reduced by many IGRs, such as the Mediterranean flour moth *Ephesia kuehniella* by tebufenozide (Khebbab *et al.*, 2008)^[46]; the European grapevine moth *Lobesia botrana* (Saenz-de-Cabezón *et al.*, 2005)^[47] and the tobacco cutworm *Spodoptera litura* (Shahout *et al.*, 2011)^[48] by methoxyfenozide; the Colorado potato beetle *Leptinotarsa decemlineata* (Farinos *et al.*, 1999)^[49] and the mealworm beetle *Tenebrio molitor* (Taibi *et al.*, 2003)^[50] by the halofenozide; *S. litura* by chlorfluazuron (Perveen and Miyata, 2000)^[51], the house fly *Musca domestica* by lufenuron (Hamadah, 2003)^[52], the red cotton stainer *Dysdercus koenigii* by flufenoxuron (Khan and Qamar, 2011)^[53]; *E. kuehniella* by diflubenzuron and hexaflumuron (Ashouri *et al.*, 2014)^[54]; the diamondback moth *Plutella xylostella* by pyriproxyfen (Mahmoudv and *et al.*, 2015)^[55]; the red flour beetle *Tribolium castaneum* (Gado *et al.*, 2015)^[56] and the onion fly *Delia antiqua* (Zhou *et al.*, 2016)^[57] by lufenuron; the rice moth *Corcyra cephalonica* by fenoxycarb (Begum and Qamar, 2016)^[58]; *P. unionalis* by Methoxyfenozide (Hamadah *et al.*, 2017)^[37] and *P. gossypiella* by Noviflumuron or Novaluron (Tanani and Ghoneim, 2018)^[38]. Results of the present study were in accordance with these reported results because treatment of penultimate instar larvae of *S. littoralis* with novaluron resulted in drastically reduced fecundity in a dose-dependent manner. A similar suppressing action on fecundity was exerted after treatment of last instar larvae with this IGR.

In contrast, the recorded result in the current study disagreed with the failure of some IGRs to halt the fecundity of some insects, such as fenoxycarb against the honey bee *Apis mellifera* (Thompson *et al.*, 2005)^[59], methoxyfenozide against the beet armyworm *Spodoptera exigua* (Christian-Lius and Pineda, 2010)^[60] and Novaluron and Diflubenzuron against the brown marmorated stink bug *Halyomorpha halys* (Kammaing *et al.*, 2012)^[61]. Furthermore, feeding of larvae on leaves treated with methoxyfenozide enhanced the fecundity of *S. littoralis* (Ishaaya *et al.*, 1995)^[62]. These diverse effects could be attributed to the different modes of action of IGRs, different insect susceptibilities, time of

treatment and some of other factors.

The drastic reduction of fecundity in *S. littoralis*, after larval treatment with novaluron in the present study, might be due to its interference with some processes, from the development of ovarian follicle to egg maturation. However, some scenarios can be discussed in this context. First, Novaluron might induce some disorders in the developing ovarioles during the immature stages (Davey, 1993)^[63], including cell death in the germarium, oocyte resorption in the pre-vitelarium and vitellarium before oviposition (Zhou *et al.*, 2016)^[57], and proliferation of follicle cells (Lucantoni *et al.*, 2006; Khan *et al.*, 2007)^[64, 65]. Second, Novaluron might inhibit the ovarian growth during the oogenesis (Smagghe *et al.*, 1996; Salem *et al.*, 1997)^[31, 66]. Third, it might disturb the production and/or function of the gonadotropic hormone responsible for the synthesis of vitellogenins and regulation of vitellogenesis (Di Ilio *et al.*, 1999)^[67]. Also, it might disturb the ecdysone activity, threshold of which is crucial for the normal oogenesis (Terashima *et al.*, 2005)^[68]. Fourth, eggs may develop normally in ovaries, but they could not be lay, because of the seriously deformed ovipositors of females or to the reduced mechanical strength (Moreno *et al.*, 1994)^[69]. Fifth, the prohibited fecundity of *S. littoralis*, in the current study, might be due to an inhibitory effect of the tested IGR on synthesis of both DNA and RNA, suboptimal nutrition because of reduced feeding, altered copulation behaviour as a result of sublethal intoxication against the tested compound, or a combination of factors.

Inhibited fertility: Fertility is a principal parameter of the reproductive capacity in insects. In the current study, novaluron exhibited a potent reducing effect on fertility of *S. littoralis*, regardless the concentration or the time of larval treatment. To a great extent, this result was in agreement with many reported results of drastically reduced fertility of the same pest by several IGRs, such as chlorfluazuron (Sammour *et al.*, 2008)^[70], methoxyfenozide (Pineda *et al.*, 2009)^[43], diflubenzuron (Aref *et al.*, 2010)^[42], lufenuron (Gaaboub *et al.*, 2012)^[45], triflumuron (El-Naggar, 2013)^[71], as well as remarkably reduced fertility of *P. gossypiella* after treatment of 1-day old eggs with hexaflumuron (El-Barkey *et al.*, 2009)^[72], lufenuron, chlorfluazuron or chromafenozide (Kandil *et al.*, 2012)^[15], chlorfluazuron or hexaflumuron (Kandil *et al.*, 2013)^[73], chromafenozide or diflubenzuron (Salem, 2015)^[74], noviflumuron or novaluron (Tanani and Ghoneim, 2018)^[38], or after treatment of newly hatched larvae with novaluron (Hassan *et al.*, 2017)^[31]. Moreover, the present result was in accordance, to some extent, with several reported results of inhibited fertility of other insect species by various IGRs, such as *S. litura* (Perveen and Miyata, 2000)^[51], *M. domestica* by diufenolan (Hamadah, 2003)^[52], *T. molitor* by Halofenozide (Taibi *et al.*, 2003)^[50], *T. castaneum* by novaluron (Kostyukovsky and Trostanelsky, 2004)^[75], *D. koenigi* by flufenoxuron (Khan and Qamar, 2011)^[53], *C. maculatus* by cyromazine (Al-Mekhlafi *et al.*, 2011)^[36], *E. kuehniella* by tebufenozide (Khebbab *et al.*, 2008)^[46], diflubenzuron or hexaflumuron (Ashouri *et al.*, 2014)^[54], methoxyfenozide (Bouzera and Soltani-Mazouni, 2014)^[76], *S. exigua* (Christian-Lius and Pineda, 2010)^[60], *S. litura* (Shahout *et al.*, 2011)^[48], by methoxyfenozide as well as *P. unionalis* methoxyfenozide (Hamadah *et al.*, 2017)^[37].

It is well known that the egg maturation in insects depends basically on the vitellogenins, including proteins, carbohydrates and lipids, all of which are prerequisite

metabolites for the embryonic development (Soltani and Mazouni, 1992; Chapman, 1998)^[77, 78]. These materials are synthesized principally by fat body during the immature stages (Telfer, 2009)^[79] or by the ovary *in situ* (Indrasith *et al.*, 1988)^[80]. For explicating the reduced fertility of *S. littoralis*, after larval treatment with novaluron, in the present investigation, it might disturb the production and/or accumulation of vitellogenins in the adult females leading to reduction of fertility (for some detail, see Taibi *et al.*, 2003; Pineda *et al.*, 2006; Osorio *et al.*, 2008)^[50, 81, 82]. On the other hand, the tested IGR might indirectly affect the fertility *via* its adverse action on "potency" of intracellular spaces in the follicular epithelium or generally prohibited the role of gonadotropic hormone responsible for the vitellogenin deposition into oocytes (Davey and Gordon, 1996)^[83]. Also, the fertility reduction might be due to the penetration of novaluron residues through mothers into eggs and disturbed the synthesis of embryonic cuticle. So, the fully mature embryos had poorly chitinized mouth parts which were insufficiently rigid to perforate the surrounding vitellin membrane for egg hatching (Marco and Vinuela, 1994; Sallam, 1999; Sammour *et al.*, 2008)^[84, 85, 70]. Another suggestion for the fertility reduction after treatment of *S. littoralis* larvae with novaluron, in the current study, can be provided herein. This compound might adversely affect the survival of developing embryos at certain stages resulting in decreasing hatching percentage. On the basis of some molecular studies, some IGRs affect the insect reproduction *via* the perturbation of gene expression in the hierarchy cascade of vitellogenesis and/or choriogenesis (Sun *et al.*, 2003)^[86], the tested IGR might interfere with the gene expression resulting in a reduction of the developing embryos in *S. littoralis*.

3. Retarded embryogenesis of *S. littoralis* by Novaluron

In insects, the egg incubation period usually indicates the rate of embryonic developmental, i.e., shorter period usually denotes faster rate and *vice versa*. In the present study, treatment of *S. littoralis* larvae with novaluron resulted generally in retarded embryonic development; since the incubation period was significantly prolonged, regardless the concentration or time of treatment. This result corroborated with little reported results recording a similar retarding action of some IGRs on the embryonic development of *P. gossypiella*, after treatment of eggs with hexaflumuron (El-Barkey *et al.*, 2009)^[72] or noviflumuron (Tanani and Ghoneim, 2018)^[38], and after treatment of newly hatched larvae with LC₅₀ values of lufenuron, chlorfluazuron or chromafenozide (Kandil *et al.*, 2012)^[15], novaluron (Hassan *et al.*, 2017)^[31]. Also, such period was prolonged after treatment of *C. maculatus* larvae with cyromazine (Al-Mekhlafi *et al.*, 2011)^[36] and after treatment of *P. unionalis* last instar larvae with methoxyfenozide (Hamadah *et al.*, 2017)^[37]. The retarded embryonic development in *S. littoralis* after treatment of 5th or 6th instar larvae with novaluron, in the present study, might be attributed to its effect on the ecdysteroids responsible for the regulation of certain stages of embryogenesis, especially those ecdysteroids originating from the ovary *in situ* (Chapman, 1998)^[78].

Conclusion

In the light of the present results, novaluron acts as an antigonadotropic compound in *S. littoralis*, reducing the

female fecundity, prohibiting the fertility and retarding the embryonic development. Therefore, this IGR can be used as an effective agent in the integrated pest management of this dangerous pest because it has developed resistance to the majority of conventional insecticides.

References

- Hill DS. Agricultural insect pests of temperate regions and their control. Cambridge University Press, 1987.
- Lal OP, Naji AH. A contribution to the host range of the vegetable armyworm *Spodoptera littoralis* (Boisd) (Lepidoptera: Noctuidae) in the Socialist Peoples, Arab Jamahriyya. J. Entomol. Res. 1990; 14(12):150-155.
- El Aswad AF, Abdelgaleil SAM, Nakatani M. Feeding deterrent and growth inhibitory properties of limonoids from *Khaya senegalensis* against the cotton leafworm, *Spodoptera littoralis*. Pest Manage. Sci. 2003; 60:199-203.
- Roques A, Rabitsch W, Rasplus JY, Lopez Vaamonde C, Nentwig W. Alien terrestrial invertebrates of Europe. In: "Handbook of alien species in Europe: Invading Nature" (Daisie, ed.). Springer Series in Invasion Ecology, Springer-Verlag, 2008, 63-80.
- Smaghe G, Carton B, Wesemael W, Ishaaya I, Tirry L. Ecdysone agonists-mechanism of action and application on *Spodoptera* species. Pestic. Sci. 1999; 55:343-389.
- Miles M, Lysandrou M. Evidence for negative cross resistance to insecticides in field collected *Spodoptera littoralis* (Boisd.) from Lebanon in laboratory bioassays. Mededelingen (Rijksuniversiteitte Gent. Fakulteit van de Landbouw kundige en Toegepaste Biologische Wetenschappen). 2002; 67:665.
- Aydin MH, Gurkan MO. The efficacy of spinosadon different strains of *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae). Tur. J. Biol. 2006; 30:5-9.
- Hussain A. The effect of non-host plant volatiles on the reproductive behaviour of the Egyptian cotton leafworm, *Spodoptera littoralis*. M.Sc. Thesis, Swedish Univ. Agric.Sci., Sweden, 2012.
- Raslan SAA. Preliminary report on initial and residual mortality of the natural product, Spinosad, for controlling cotton leaf worm egg masses. In: Egypt. 2nd Inter. Conf., Plant Prot. Res. Inst., Cairo, Egypt, 21-24 December. 2002; 1: 635-637.
- Gurr GM, Thwaite WG, Nicol HI. Field evaluation of the effects of the insect growth regulator tebufenozide on entomophagous arthropods and pests of apples. Australian J. Entomol. 1999; 38:135-140.
- Tunaz H, Uygun N. Insect growth regulators for insect pest control. Turkish J Agric. Forestry. 2004; 28:337-387.
- Hammock CD, Quistad GB. Meabolism and mode of action of juvenile hormone, juvenoids and other insect growth regulators. In: "Progress in pesticide Biochemistry" (Hutson DH, Roberts TR. eds.). John Wiley & Sons Ltd. 1981; 1:1-85.
- Ishaaya I, Horowitz AR, Tirry L, Barazani A. Novaluron (Rimon), a novel IGR: mechanism, selectivity and importance in IPM programs. Proc. Int. Symp. Crop Protect. Med. Fac. Landbouww Univ., Gent. 2002; 67: 617-626.
- Ishaaya I, Kontsedalov S, Horowitz AR. Novaluron (Rimon), a novel IGR: Potency and cross-resistance. Arch. Insect Bioch. Physiol. 2003; 54:157-164.
- Kandil MA, Ahmed AF, Moustafa HZ. Toxicological and biochemical studies of lufenuron, chlorfluazuron and chromafenozide against *Pectinophora gossypiella* (Saunders). Egypt. Acad. J. Biolog. Sci. 2012; 4(1):37-47.
- Soltani N, Besson MT, Delachambre J. Effects of diflubenzuron on the pupal-adult development of *Tenebrio molitor* (L.) (Coleoptera: Tenebrionidae): growth and development cuticle secretion, epidermal cell density, and DNA synthesis. Pestic. Biochem. Physiol. 1984; 21:256-264.
- Cutler GC, Scott Dupree CD, Tolman JH, Harris CR. Field efficacy of novaluron for control of Colorado potato beetle (Coleoptera: Chrysomelidae) on potato. Crop Prot. 2007; 26:760-767.
- Alyokhin A, Guillemette R, Choban R. Stimulatory and suppressive effects of Novaluron on the Colorado potato beetle reproduction. J. Econ. Entomol. 2009; 102(6):2078-2083.
- Kim KS, Chung BJ, Kim HK. DBI-3204: A new benzoylphenyl urea insecticide with a particular activity against whitefly. Brighton Crop Prot. Conf. 2000; 1:41-46.
- Ishaaya I, Kontsedalov S, Masirov D, Horowitz AR. Biorational agents-mechanism, selectivity and importance in IPM programs for controlling agricultural pests. Med. Landbouww Rijksuniv Gent. 2001; 66:363-374.
- Barazani A. Rimon, an IGR insecticide. Phytoparasitica. 2001; 29:59-60.
- Ishaaya I, Horowitz AR. Novaluron (Rimon) a novel IGR: its biological activity and importance in IPM programs. Phytoparasitica. 2002; 30:203.
- Malhata FM, Loutfy NM, Ahmed MT. Dissipation kinetics of novaluron in tomato, an arid ecosystem pilot study. Toxic. Environ. Chem. 2014; 96(1):41-47.
- Ghoneim, KS. Physiological studies on endocrine and reproductive systems of the cotton leafworm *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). Ph.D. Thesis, Fac. of Sci., Al-Azhar Univ., Cairo, Egypt, 1985.
- Bakr RFA, El-barky NM, Abd Elaziz MF, Awad MH, Abd El-Halim HME. Effect of Chitin synthesis inhibitors (flufenoxuron) on some biological and biochemical aspects of the cotton leaf worm *Spodoptera littoralis* Bosid. (Lepidoptera: Noctuidae). Egypt. Acad. J Biolog. Sci. 2010; 2(2):43-56.
- Topozada A, Abd-allah S, El-Defrawi ME. Chemosterilization of larvae and adults of the Egyptian cotton leafworm, *Prodenia litura* by apholate, metepa and hempa. J. Econ. Entomol. 1966; 59:1125-1128.
- Moroney MJ. Facts from figures (3rd ed.). Penguin Books Ltd., Harmonds worth. Middle Sex, 1956.
- Williams D, Amos TG. Some effects of synthetic juvenile hormones and hormone analogues on *Tribolium castaneum* (Herbs"). Aust. J Zool. 1974; 22:147-53.
- Ghoneim K, Al-Daly A, Amer M, Mohammad A, Khadrawy F, Mahmoud MA. Disruptive effects of pomegranate *Punica granatum* Linn. (Lythraceae) extracts on the reproductive potential of desert locust *Schistocerca gregaria* (Forsk.) (Orthoptera: Acrididae). Entomology and Applied Science Letters. 2014; 1(1):1-10.
- Mondal KAMSH, Parween S. Insect growth regulators and their potential in the management of stored product

- insect pests. *Integ. Pest Manage. Rev.* 2000; 5:255-295.
31. Hassan HA, Ghoneim K, Tanani MA, Bakr NA. Impaired adult performance and reproductive potential of the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) by the chitin synthesis inhibitor, Novaluron. *Journal of Entomology and Zoology Studies.* 2017; 5(2):581-592.
 32. Bakr RFA, Guneidy NAM, El-Bermawy SM. Toxicological and biological activities of three IGRs against fourth larval instar of the cotton leafworm *Spodoptera littoralis* (Boisd.). *J. Egypt. Acad. Environ. Develop.* 2005; 6(4):103-132.
 33. Soltani-Mazouni N, Soltani N. Diflubenzuron affected DNA synthesis in the ovaries of *Tenebrio molitor*. *J. Invert. Reprod. Devel.* 1994; 25:19-21.
 34. Al Dali AG, Ghoneim KS, Bakr RF, Bream AS, Tanani MA. Egg productivity of *Schistocerca gregaria* (Orthoptera: Acrididae) as affected by the non-steroidal ecdysone agonist Tebufenozide (RH-5992). *J. Egypt. Acad. Soc. Environ. Develop.* 2008; 9(10):27-38.
 35. Smagghe G, Degheele D. Action of a novel non-steroidal ecdysteroid mimics, tebufenozide (RH-5992), on insects of different orders. *Pestic. Sci.* 1994; 42:85-92.
 36. Al Mekhlafi F, Mashaly AM, Abdel Mageed A, Wadaan MA, Al-Mallah NM. Overlap effects of cyromazine concentration, treatment method and rearing temperature on the Southern cowpea weevil (*Callosobruchus maculatus* F.) reared on cowpea. *Afr. J. Microbio. Res.* 2011; 5(32):5848-5853.
 37. Hamadah Kh, Ghoneim K, Mansour AN, Abo Elsoud AA. Deranged adult performance and reproductive potential of the olive leaf moth *Palpita unionalis* (Hübner)(Lepidoptera: Pyralidae) by the Non-steroidal ecdysone agonist, Methoxyfenozide. *International Journal of Information Research and Review.* 2017; 04(06):4228-4240.
 38. Tanani M, Ghoneim K. Disruptive effects of certain chitin synthesis inhibitors on adult performance and reproductive potential of the pink bollworm, *Pectinophora gossypiella* (Saunders)(Lepidoptera: Gelechiidae). *Egyptian Academic Journal of Biological Sciences.* 2018; 11(5):79-102.
 39. Behrens W, Hoffman KH. Effects of exogenous ecdysteroids on reproduction in cricket *Gryllus bimaculatus*. *Int. Invert. Reprod. Develop.* 1983; 6:149-159.
 40. Smagghe G, Vinuela E, Budia F, Degheele D. *In vivo* and *in vitro* effects of the nonsteroidal ecdysteroid agonist tebufenozide on cuticle formation in *Spodoptera exigua*: an ultrastructural approach. *Arch. Insect Biochem. Physiol.* 1996; 32:121-134.
 41. Parween S, Faruki SI, Begum M. Impairment of reproduction in the red flour beetle, *Tribolium castaneum* (Herbst.)(Coleoptera: Tenebrionidae) due to larval feeding on triflumuron-treated diet. *J. Appl. Entomol.* 2001; 125:1-4.
 42. Aref SA, Bayoumi OCh, Soliman HAB. Effect of certain insecticides on the biotic potential of the cotton leafworm, *Spodoptera littoralis* (Boisd.). *Egypt. J. Agric. Res.* 2010; 88(1):31-40.
 43. Pineda S, Martinez AM, Figueroa JI, Schneider MI, Del Estal P, Vinuela E, Gomez B, Smagghe G, Budia F. Influence of azadirachtin and methoxyfenozide on life parameters of *Spodoptera littoralis*. *J. Econ. Entomol.* 2009; 102:1490-1496.
 44. Abdel Rahman SM, Hegazy EM, Elweg AE. Direct and latent effect of two chitin synthesis inhibitors to *Spodoptera littoralis* larvae (Boisd.). *American Eurasian J. Agric. Environ. Sci.* 2007; 2(4):454-464.
 45. Gaaboub I, Halawa S, Rabiha A. Toxicity and biological effects of some insecticides, IGRs and Jojoba oil on Cotton Leafworm *Spodoptera littoralis* (Boisd.). *J. App. Sci. Res.* 2012; 8(10):51-61.
 46. Khebbeb MEH, Gaouaoui R, Bendjeddou F. Tebufenozide effects on the reproductive potentials of the Mediterranean flour moth, *Ephesia kuehniella*. *Afr. J Biotech.* 2008; 7(8):1166-1170.
 47. Saenz de Cabezon IFJ, Marco V, Salmo FG, Perez-Moreno I. Effects of methoxyfenozide on *Lobesia botrana* Den and Schiff (Lepidoptera: Tortricidae) egg, larval and adult stages. *Pest Manage. Sci.* 2005; 11:1133-1137.
 48. Shahout HA, Xu J-X, Qiao J, Jia Q-D. Sublethal effects of Methoxyfenozide, in comparison to Chlorfluazuron and Beta-Cypermethrin, on the reproductive characteristics of common cutworm *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae). *J. Entomol. Res. Soc.* 2011; 13(3):53-63.
 49. Farinos GP, Smagghe G, Tirry L, Castañera P. Action and pharmacokinetics of a novel insect growth regulator, halofenozide, in adult beetles of *Aubeonymus mariaefrancisciae* and *Leptinotarsa decemlineata*. *Arch. Insect Biochem. Physiol.* 1999; 41:201-213.
 50. Taibi F, Smagghe G, Amrani L, Soltani-Mazouni N. Effect of ecdysone agonist RH-0345 on reproduction of mealworm, *Tenebrio molitor*. *Comp. Biochem. Physiol.* 2003; 135:257-267.
 51. Perveen F, Miyata T. Effects of sublethal dose of Chlorfluazuron on ovarian development and oogenesis in the common cutworm *Spodoptera litura* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Amer.* 2000; 93(5):1131-1137.
 52. Hamadah KhSh. Physiological and Biochemical Effects of IGRs and plant extracts on the house fly *Musca domestica*. M.Sc. Thesis, Fac. Sci., Al-Azhar Univ., Cairo, Egypt, 2003.
 53. Khan I, Qamar A. Biological activity of andalin (flucycloxuron), a novel chitin synthesis inhibitor, on red cotton stainer *Dysdercus koenigii* (Fabricius). *Frontiers in Life Sciences: Basic and Applied Biology and Medicine.* 2011; 3(2):324-335.
 54. Ashouri S, Pourabad RF, Ebadollahi A. The effect of diflubenzuron and hexaflumuron on the last larval instars of the Mediterranean flour moth *Anagasta kuehniella* (Zeller) (Lepidoptera: Pyralidae) under laboratory conditions. *Archives of Phytopathology and Plant Protection.* 2014; 47(1):75-81
 55. Mahmoudvand M, Moharramipour S, Iranshahi M. Effects of pyriproxyfen on life table indices of *Plutella xylostella* in multigenerations. *Psyche*, 2015. Article ID 453701, 7 pp. <http://dx.doi.org/10.1155/2015/453701>
 56. Gado P, Salokhe SG, Deshpande SG. Impact of Lufenuron (5.4% EC) on reproductive end points of *Tribolium castaneum*. *World Journal of Pharmaceutical Research.* 2015; 4(3):1593-1599.
 57. Zhou F, Zhu G, Zhao H, Wang Z, Xue M, Li X, Xu H, Ma X, Liu Y. Sterilization effects of adult-targeted baits containing insect growth regulators on *Delia antiqua*.

- Sci. Rep. 2016; 6:32855. 9pp. doi: 10.1038/srep32855.
58. Begum R, Qamar A. Fenoxycarb- a potent inhibitor of metamorphosis and reproduction in Rice Moth, *Corcyra cephalonica* (Stainton). Journal of Entomology and Zoology Studies. 2016; 4(4):572-577.
 59. Thompson HM, Wilkins S, Battersby AH, Waite RJ, Wilkinson D. The effects of four insect growth-regulating (IGR) insecticides on honeybee (*Apis mellifera* L.) colony development, queen rearing and drone sperm production, Ecotoxicology. 2005; 14:757-769.
 60. Christian Luis R, Pineda S. Toxicity and sublethal effects of methoxyfenozide on *Spodoptera exigua* (Lepidoptera: Noctuidae). J. Econ. Entomol. 2010; 103:662-667.
 61. Kamminga KL, Kuhar TP, Wimer A, Herbert DA. Effects of the insect growth regulators novaluron and diflubenzuron on the brown marmorated stink bug. Plant Health Progress Online, 2012. doi:10.1094/PHP-2012-1212-01-RS.
 62. Ishaaya I, Yablonski S, Horowitz AR. Comparative toxicity of two ecdysteroid agonists, RH-2485 and RH-5992, on susceptible and pyrethroid-resistant strains of the Egyptian cotton leafworm, *Spodoptera littoralis*. Phytoparasitica. 1995; 23:139-145.
 63. Davey KG. Hormonal integration of egg production in *Rhodnius prolixus*. Am. Zool. 1993; 33:397-402.
 64. Lucantoni L, Giusti F, Cristofaro M, Pasqualini L, Esposito F, Lupetti P, Habluetzel A. Effects of a neem extract on blood feeding, oviposition and oocyte ultrastructure in *Anopheles stephensi* Liston (Diptera: Culicidae). Tissue Cell. 2006; 38:361-371.
 65. Khan M, Hossain MA, Islam MS. Effects of neem leaf dust and a commercial formulation of a neem compound on the longevity, fecundity and ovarian development of the melon fly, *Bactrocera cucurbitae* (Coquillett) and the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). Pak. J. Biol. Sci. 2007; 10:3656-3661.
 66. Salem H, Smagghe G, Degheele D. Effects of tebufenozide on oocyte growth in *Plodia interpunctella*. Medical. Faculty. Landbouww. Gent University. 1997; 62(1):9-13.
 67. Di Ilio V, Cristofaro M, Marchini D, Nobili P, Dallai R. Effects of a neem compound on the fecundity and longevity of *Ceratitidis capitata* (Diptera: Tephritidae). J. Econ. Entomol. 1999; 92:76-82.
 68. Terashima J, Takaki K, Sakurai S, Bownes M. Nutritional status affects 20-hydroxyecdysone concentration and progression of oogenesis in *Drosophila melanogaster*. J. Endocrinol. 2005; 187:69-79.
 69. Moreno DS, Martinez AJ, Riviello MS. Cyromazine effects on the reproduction of *Anastrepha ludens* (Diptera: Tephritidae) in the laboratory and in the field. J. Econ. Entomol. 1994; 87:202-211.
 70. Sammour EA, Kandil MA, Abdel-Aziz NF. The reproductive potential and fate of chlorfluazuron and lufenuron against cotton leafworm, *Spodoptera littoralis* (Boisd.). American Eurasian J. Agric. Environ. Sci. 2008; 4(1):62-67.
 71. El Naggar J. Sublethal effect of certain insecticides on biological and physiological aspects of *Spodoptera littoralis* (Boisd.). Nature & Science. 2013; 11(7):19.
 72. El Barkey NM, Amer AE, Kandil MA. Ovicidal activity and biological effect of Radiant and hexaflumuron against eggs of pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). Egypt. Acad. J. Biol. Sci. 2009; 2:23-36.
 73. Kandil MAA, Salem MS, Adly AM. Biological and biochemical changes in pink bollworm, *Pectinophora gossypiella* after treatment with Hexaflumuron and Chlorfluazuron. Annals of Agric. Sci., Moshtohor (Egypt). 2013; 51(4):472-437.
 74. Salem MSM. Latent effect of different compounds on *Pectinophora gossypiella* (Saunders). J. Plant Prot. and Path., Mansoura Univ., Egypt. 2015; 6(2):269-279.
 75. Kostyukovsky M, Trostanetsky A. The effect of a new chitin synthesis inhibitor, Novaluron, on various developmental stages of *Tribolium castaneum* (Herbst). J. stored Product Res. 2004; 42(2):136-148.
 76. Bouzera H, Soltani-Mazouni N. Comparative effects of two moulting hormone agonists (Methoxyfenozide and Tebufenozide) on the Mediterranean flour moth *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae): ecdysteroids amounts of testes and reproductive events. World Appl. Sc. J. 2014; 31 (11):1903-1910.
 77. Soltani N, Mazouni A. Diflubenzuron and oogenesis in the codling moth, *Cydia pomonella*. Pesti. Sci. 1992; 34:257-261.
 78. Chapman RF. The insects: structure and Function. 4thed. Cambridge: Cambridge University Press 1998, 116-118.
 79. Telfer WH. Egg formation in Lepidoptera. J. Insect Sci. 2009; 9:50. (insectscience.org/9.50).
 80. Indrasith L, Sasaki ST, Yaginuma T, Yamashita O. The occurrence of premature form of egg-specific protein in vitellogenic follicles of *Bombyx mori*. J.Comp. Physiol. 1988; 158:1-7.
 81. Pineda S, Smagghe G, Schneider MI, Del Estal P, Vinuela E, Martinez AM, Budia F. Toxicity and pharmacokinetics of spinosad and methoxyfenozide to *Spodoptera littoralis* (Lepidoptera: Noctuidae). Environ. Entomol. 2006; 35:856-864.
 82. Osorio A, Martinez AM, Schneider MI, Diaz O, Corrales JL, Aviles MC, Smagghe G, Pineda S. Monitoring of beet army worm resistance to spinosad and methoxyfenozide in Mexico. Pest Manag. Sci. 2008; 64:1001-1007.
 83. Davey KG, Gordon DRB. Fenoxycarb and thyroid hormones have JH-like effects on the follicle cells of *Locosta migratoria* in vitro. Arc Insect Biochem.Physiol. 1996; 32:613- 626.
 84. Marco V, Vinuela E. Effects of hexaflumuron on fecundity, fertility and longevity of *Ephestia kuehniella* Zeiler and *Spodoptera exigua* (Hubner). Med. Fac. Landbouww Umv. Gent, 1994, 59/2a.
 85. Sallam MH. Effect of Diflubenzuron on embryonic development of the acridid, *Heteracris littoralis*. J. Egypt. Ger. Soc. Zool. 1999; 30(E):17-26.
 86. Sun X, Song Q, Barrett B. Effect of ecdysone agonists on vitellogenesis and the expression of EcR and USP in codling moth (*Cydia pomonella*). Arch. Insect Biochem. Physiol. 2003; 52:115-129.