



Biochemical profiling of the cestode *Moniezia* sp. and its host *Capra hircus* in the Nanded region of Maharashtra

Kadam Pooja Mahajan¹, Dhanraj Balbhim Bhure^{2*}

¹ Research Scholar, Department of Zoology, Yeshwant Mahavidyalaya, Nanded, Maharashtra, India

² Research Supervisor, Department of Zoology, Yeshwant Mahavidyalaya, Nanded, Maharashtra, India

Corresponding Author: Dhanraj Balbhim Bhure

DOI: <https://doi.org/10.66856/ijbr.2026.11.2.11035>

Abstract

The present investigation was undertaken to evaluate the biochemical composition of the cestode parasite *Moniezia* sp. and the intestinal tissue of its host, *Capra hircus*. The concentrations of major biomolecules, namely protein, glycogen, and lipid, were estimated using standard biochemical methods. Protein content was recorded as 16.15 mg/g in the normal intestine, 13.17 mg/g in the infected intestine, and 12.64 mg/g in *Moniezia* sp. Glycogen content was 21.36 mg/100 ml in the normal intestine, 18.15 mg/100 ml in the infected intestine, and 16.38 mg/100 ml in the parasite. Similarly, lipid content was found to be 15.72 mg/g in the normal intestine, 12.57 mg/g in the infected intestine, and 11.78 mg/g in the parasite. A progressive decline in all biochemical constituents was observed from normal host tissue to infected tissue and further to the parasite. These findings indicate that *Moniezia* sp. depends largely on host-derived nutrients and causes significant biochemical alterations in the host intestine. The study provides valuable information regarding host–parasite metabolic interactions and contributes baseline data on the biochemical ecology of cestode infections in goats from the Nanded region of Maharashtra.

Keywords: *Moniezia* sp., *Capra hircus*, Biochemistry, Protein, Glycogen, Lipid, Host–parasite interaction

Introduction

Goats (*Capra hircus*) make up 26.40% of India's livestock, establishing it as a leading goat-rearing nation (Anonymous, 2012) [1]. Goat farming is crucial for the rural economy, offering income, nutrition, and livelihood for marginal farmers, landless workers, and nomadic groups (Singh, *et al.*, 2018; Deshmukh and Shinde, 2020) [6, 30]. Gastrointestinal parasitic infections, especially cestodes, significantly reduce goat productivity (Patel, *et al.*, 2019) [22]. Cestodes absorb nutrients from the host's intestine, causing malnutrition, stunted growth, anemia, and reduced productivity, impacting the health and economic value of animals (Gaikwad *et al.*, 2017; Pawar and Hiware, 2020; Shinde *et al.*, 2018; Bhosale *et al.*, 2021) [3, 9, 24, 28]. Biochemistry, which studies biomolecules' composition and functions, offers insights into parasite physiology and host–parasite interactions, elucidating metabolic pathways and survival strategies of parasites. Cestodes, relying on host nutrients, are ideal for studying physiological and biochemical processes (Fairbairn *et al.*, 1961; Smyth and McManus, 1989; Barratt, 1981) [7, 31]. They depend on glucose metabolism for energy, storing carbohydrates like glycogen as reserves (Mishra *et al.*, 1945-1991; Markov, 1943; Read and Rothman, 1957; Kulkarni and Patil, 2017) [16, 20, 21, 26]. Proteins, comprising 20–40% of cestodes' dry weight, are crucial for structure, enzymatic activity, and metabolic regulation (Barrett, 1981; Shaikh, *et al.*, 2020) [27]. Lipids, vital for reproduction and energy storage, are abundant in cestodes, which rely on host nutrients (Brand and Van T., 1952). The genus *Moniezia*, in the family Anoplocephalidae, commonly infects grazing ruminants like goats and sheep in tropical and subtropical regions. Several *Moniezia* species in Indian ruminants show taxonomic diversity and ecological adaptation, with many documented

in Maharashtra's *Capra hircus*, highlighting cestode biodiversity (Jadhav and Ghogare, 2016; Mane *et al.*, 2019) [12, 19].

Despite the significance of cestode infections, biochemical studies on cestodes in India remain limited. Previous investigations in Maharashtra's Marathwada region have documented significant variations in protein, glycogen, and lipid content among cestode parasites, reflecting adaptive metabolic strategies influenced by host nutritional status, parasite developmental stage, and environmental factors (Chavan *et al.*, 2021; Ghorband and Pawar, 2018; Patil *et al.*, 2021) [5, 10, 23]. These findings underscore the importance of biochemical profiling in understanding parasite physiology, host exploitation strategies, and potential therapeutic targets. However, no comprehensive biochemical investigation of *Moniezia* from the Nanded region of Maharashtra has been conducted. This study aims to evaluate the biochemical composition of *Moniezia* and its impact on the infected intestine of *Capra hircus* from the Nanded region of Maharashtra. The outcomes are expected to provide valuable insights into cestode metabolism, host–parasite relationships, and regional parasitic ecology, contributing meaningful baseline data for future research in parasitology, biochemistry, and disease management strategies.

Material and Methods

Collection of Material

The worms were removed from *Capra hircus* digestive tract and cleaned with distilled water. Collected worms were dried on blotting paper to eliminate excess water before being transferred to a watch glass and weighed using a

sensitive balance. The dry weight was also determined after 24 hours of 50-60 Co exposure.

Biochemical Estimation

The biochemical estimation followed by the following standard methods.

1. The estimation of protein content in the cestode parasites were carried out by Lowry's method (1951)^[17]

2. The glycogen estimation was carried out by Kemp *et al.*, (1954)^[15] method.
3. The lipid estimation by Folch *et al.*, (1957)^[8] method.

Result and Discussion

The biochemical composition of the normal intestine, infected intestine of *Capra hircus*, and the cestode parasite *Moniezia* sp. is presented in Table 1 and Figure 1.

Table 1: Biochemical estimation of ruminant *Capra hircus* intestine and cestode i.e. *Moniezia*

Name of parameter	Host intestine (<i>Capra hircus</i>)	Host infected intestine (<i>Capra hircus</i>)	Cestode Parasite (<i>Moniezia</i>)
Protein	16.15 mg/ gm. wt. of tissue	13.17 mg/gm. wt. of tissue	12.64 mg/gm. wt. of tissue
Glycogen	21.36 mg/100 ml of solution	18.15 mg/100 ml of solution	16.38 mg/100 ml of solution
Lipid	15.72 mg/gm. of wt. of tissue	12.57 mg/gm. of wt. of tissue	11.78 mg/gm. of wt. of tissue

Protein content was highest in the normal intestine (16.15 mg/g tissue), followed by the infected intestine (13.17 mg/g tissue) and *Moniezia* sp. (12.64 mg/g tissue). The infected intestine showed an 18.45% reduction, whereas the parasite exhibited a 21.73% reduction relative to the normal intestine.

Glycogen content was recorded as 21.36 mg/100 ml in the normal intestine, 18.15 mg/100 ml in the infected intestine, and 16.38 mg/100 ml in *Moniezia* sp. The reduction was

15.03% in the infected intestine and 23.31% in the parasite when compared with normal tissue.

Similarly, lipid content was highest in the normal intestine (15.72 mg/g tissue), followed by the infected intestine (12.57 mg/g tissue) and the parasite (11.78 mg/g tissue). Lipid content decreased by 20.04% in infected tissue and by 25.06% in the parasite relative to the normal intestine.

The overall results indicate a progressive decrease in biochemical constituents from normal intestinal tissue to infected tissue and further to the parasite.




Fig 1: Biochemical estimation of ruminant *Capra hircus* intestine and cestode i.e. *Moniezia*

The present investigation revealed significant biochemical variations among normal intestine, infected intestine, and *Moniezia* sp. The reduction in protein, glycogen, and lipid content observed in infected intestinal tissue suggests metabolic stress imposed by parasitic infection and utilization of host nutrients by the parasite.

Protein is an essential structural and functional component involved in enzyme synthesis, tissue repair, and metabolic regulation. The observed reduction in protein content in infected tissues and the parasite may be attributed to continuous utilization of host-derived amino acids for parasite growth and maintenance. Similar observations have been reported by Ingle *et al.* (2022)^[11] and Bhogil *et al.* (2022)^[2].

Glycogen serves as the principal storage form of carbohydrate and represents an important energy reserve. The lower glycogen levels observed in infected tissues and the parasite indicate active utilization of carbohydrate reserves to meet energy requirements. Comparable findings were reported by Read and Rothman (1957)^[26], who

emphasized the importance of carbohydrate metabolism in cestode survival.

Lipids play a vital role in energy storage, membrane formation, and reproductive activities. The reduced lipid content recorded in *Moniezia* sp. and infected intestinal tissue suggests increased utilization of lipid reserves and dependence on host-derived nutrients. Similar findings have been documented by Makne (2021)^[18] and Pawar *et al.* (2022)^[25].

The gradual decline in biochemical constituents from normal intestine to infected intestine and parasite demonstrates the metabolic dependence of *Moniezia* sp. on the host. Since cestodes lack a digestive system, nutrient absorption occurs directly through the tegument, representing an efficient adaptation for survival within the host environment.

The present findings agree with previous studies on cestode parasites and support the concept that host nutrient exploitation and biochemical modulation constitute important adaptive mechanisms for parasite establishment, growth, and reproduction.

Conclusion

The present study demonstrated significant biochemical alterations associated with *Moniezia* sp. infection in *Capra hircus*. Protein, glycogen, and lipid contents were highest in normal intestinal tissue and progressively decreased in infected intestinal tissue and the parasite. These findings indicate that *Moniezia* sp. relies extensively on host-derived nutrients and exerts a considerable influence on host metabolism. The study contributes valuable baseline information on host-parasite biochemical interactions and may be useful for understanding cestode physiology and developing effective parasite management strategies in goats.

Acknowledgement

The authors are grateful to the Head, Department of Zoology, Yeshwant Mahavidyalaya, Nanded, for providing laboratory facilities necessary for the completion of this work. The authors also express their sincere thanks to the Principal, Yeshwant Mahavidyalaya, Nanded, for institutional support and encouragement throughout the study.

References

1. Anonymous. 19th Livestock census-2012. All India Report, Department of Animal Husbandry, Dairying and Fisheries, Ministry of Agriculture, Government of India, 2012.
2. Bhogil A, Borde S, Thosar A. Study of biochemical components of cestode parasites from *Capra hircus* (L.) in Solapur District (M.S.), India. *Uttar Pradesh Journal of Zoology*,2022;43(14):74–77.
3. Bhosale SR, Jadhav BV, Patil RT. Prevalence of gastrointestinal helminths in goats from Marathwada region of Maharashtra. *Journal of Parasitic Diseases*,2021;45(2):312–318. <https://doi.org/10.1007/s12639-021-01327-4>
4. Brand T, Von Chemical physiology of endoparasitic animals. Academic press, New York.
5. Chavan PB, Shinde GB, Pawar SP. Impact of gastrointestinal helminths on growth and productivity of goats in Maharashtra. *Indian Journal of Animal Sciences*,2021;91(4):301–306.
6. Deshmukh SS, Shinde RS. Role of goats in rural economy and livelihood security in Maharashtra. *Indian Journal of Extension Education*,2020;56(3):45–50.
7. Fairbairn DG, Werthim RP, Harpur, Schiller EL. Biochemistry of normal and irradiated strains of *Hymenolepis diminuta*. *Exp. Parasitol*,1961;11:248–263.
8. Folch J, Lees M, Sloane-Stanley GH. The method of lipid estimation. *J. Biol. Chem*,1957;228:497.
9. Gaikwad SS, Jadhav BV, Patil AM. Epidemiological studies of cestode infections in goats from western Maharashtra. *Uttar Pradesh Journal of Zoology*,2017;37(2):101–108.
10. Ghorband CM, Pawar SS. Biochemical alterations in cestode parasites infecting small ruminants from Marathwada region. *International Multidisciplinary Research Journal*,2018;8(2):15–21.
11. Ingle J, Thosar A, Hamde VS. Biochemical estimation of the cestode parasite, *Moniezia* sp. from *Capra hircus* (L.) in Marathwada region (M.S.), India. *Global Journal for Research Analysis*,2022;11(2):30–31.
12. Jadhav BV, Ghogare MR. Taxonomic studies on cestode parasites of domestic ruminants from Maharashtra, India. *Indian Journal of Helminthology*,2016;33(1):47–55.
13. Jadhav BV, Shinde SL. Biochemical adaptations of cestode parasites in domestic animals. *Journal of Parasitic Diseases*,2021;45(3):612–618. <https://doi.org/10.1007/s12639-020-01289-3>
14. John Barrett: Book of “Biochemistry of parasitic helminthes”, 1981.
15. Kemp A. Vankits, Haljningem AJM. A colorimetric method for the determination of glycogen in tissue. *Biochem J*, 1954, 646-648.
16. Kulkarni GK, Patil SR. Biochemical adaptations in cestode parasites of domestic animals. *Bioscience Discovery*,2017;8(3):501–507.
17. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. The method for protein estimation. *J. Biol. Chem*,1951;193:265. (The original method).
18. Makne HD. Biochemical studies of *Aliezia* tapeworm in goat *Capra hircus*. *International Journal of Zoology Studies*,2021;6(2):26–27.
19. Mane UH, Hiware CJ, Pawar RS. Diversity of cestode parasites infecting goats from Maharashtra. *Journal of Experimental Zoology India*,2019;22(2):789–796.
20. Markov GS. The dynamic of reserve nutritive substances in parasitic worms in artificial media. *Zool. Zh*,1943;22:3-18.
21. Mishra, *et al.* On a new species of the genus *Oochoristica* from the intestine of *Calotes versicolor*. *proc.Ind. Acad.Sci., section*,1945;22(17):1-5.
22. Patel PR, Singh VK, Yadav CL. Economic impact of gastrointestinal parasites in goats: An Indian perspective. *Indian Journal of Small Ruminants*,2019;25(1):10–16.
23. Patil AM, Jadhav BV, Gaikwad SS. Biochemical profiling of cestode parasites from small ruminants. *Journal of Applied and Natural Science*,2021;13(1):90–96.
24. Pawar RS, Hiware CJ. Seasonal dynamics of cestode infections in goats from drought-prone regions of Maharashtra. *Indian Journal of Veterinary Sciences and Biotechnology*,2020;15(4):42–48.
25. Pawar SD, Patil AS, Jadhav BV. Comparative biochemical studies on cestode parasites and their ruminant hosts. *Indian Journal of Parasitology*,2022;46(1):23–29.
26. Read CP, Rothman AH. The role of carbohydrates in the biology of cestodes. The effect of starvation on glycogenesis and glucose consumption in *Hymenolepis*. *Exp. Parasitol*,1957;6:280-387.
27. Shaikh AR, Pathan AA, Syed MA. Biochemical changes in helminth parasites under host nutritional stress. *Indian Journal of Fundamental and Applied Life Sciences*,2020;10(2):135–142.
28. Shinde GB, Jadhav BV, Patil RT. Prevalence and pathology of cestode infections in goats from Maharashtra. *Journal of Parasitic Diseases*,2018;42(3):367–373.
29. Shinde SL, Pawar RM, Jadhav BV. Biochemical analysis of cestode parasites in relation to their hosts. *International Journal of Zoological Studies*,2019;4(2):45–50.
30. Singh MK, Kumar R, Yadav AK. Status and prospects of goat farming in India. *Indian Journal of Animal Sciences*,2018;88(10):1201–1210.
31. Smyth JD, McManus DP. The physiology and biochemistry of Cestodes. Cambridge University Press, 1989.